

# A Note on a Line Graph of the Zero Divisor Graph of a Commutative Ring

Jaydeep Parejiya<sup>1,\*</sup>, Yesha Hathi<sup>1</sup> and Patat Sarman<sup>1</sup>

<sup>1</sup> Department of Mathematics, Government Polytechnic, Rajkot, Gujarat, India.

**Abstract:** The rings considered in this article are commutative with identity  $1 \neq 0$ . Recall that the zero divisor graph of a ring  $R$  is a simple undirected graph whose vertex set is the set of all nonzero zero divisors of the ring  $R$  and two distinct vertices  $x, y$  are adjacent in this graph if and only if  $xy = 0$ . In this article we studied the line graph of the zero divisor graph of a ring and we proved some results regarding the diameter of the line graph.

**MSC:** 13A15, 05C25.

**Keywords:** Line graph, zero divisor graph, diameter of a graph.

© JS Publication.

## 1. Introduction

The rings considered in this article are commutative with identity  $1 \neq 0$ . In 1988, Beck [6] defined the concept of zero divisor graph of a commutative ring  $R$ , where the vertices of this graph are all elements in the ring and two vertices  $x, y$  are adjacent in this graph if and only if  $xy = 0$ . Anderson and Livingston in [3] modified the definition of zero divisor graphs by restricting the vertices to the nonzero zero divisors of the ring  $R$ . The zero divisor graph was extensively studied in [1–3, 6]. The authors K. Nazzal and M. Ghanem in [8], studied the line graph of zero divisor graph. Let  $G$  be a simple undirected finite graph. Recall from [8] that line graph of  $G$  is denoted as  $L(G)$  is defined to be the graph whose vertices are the edges of  $G$ , with two vertices being adjacent if the corresponding edges share a vertex in  $G$ . This article is motivated by the interesting theorem proved on line graph of zero divisor graph of ring  $R$  in [8, 9].

It is useful to recall the following definitions from graph theory before we describe the results that are proved in this article on  $L(\Gamma(R))$ . Let  $G = (V, E)$  be a graph. Let  $a, b \in V$  with  $a \neq b$ . Recall that the *distance* between  $a$  and  $b$ , denoted by  $d(a, b)$  is defined as the length of a shortest path in  $G$  if there exists such a path in  $G$ ; otherwise, we define  $d(a, b) = \infty$ . We define  $d(a, a) = 0$ . The *diameter* of  $G$ , denoted by  $diam(G)$  is defined as  $diam(G) = \sup\{d(a, b) | a, b \in V\}$  [5]. A simple graph  $G = (V, E)$  is said to be *complete* if every pair of distinct vertices of  $G$  are adjacent in  $G$  [5, Definition 1.1.11]. Recall from [5, Definition 1.2.2], that a *clique* of  $G$  is a complete subgraph of  $G$ . A subset  $S$  of  $G$  is said to be an *independent set* if no two members of  $S$  are adjacent in  $G$ . A graph  $G = (V, E)$  is said to be *bipartite* if  $V$  can be partitioned into nonempty subsets  $V_1$  and  $V_2$  such that each edge of  $G$  has one end in  $V_1$  and the other in  $V_2$ . A bipartite graph with vertex partition  $V_1$  and  $V_2$  is said to be *complete* if each element of  $V_1$  is adjacent to every element of  $V_2$ . A complete bipartite graph with vertex partition  $V_1$  and  $V_2$  is called a *star* if either  $|V_1| = 1$  or  $|V_2| = 1$  [5, Definition 1.1.12].

\* E-mail: [parejiyajay@gmail.com](mailto:parejiyajay@gmail.com)

Let  $R$  be any ring. We denote the set of all zero divisor of ring  $R$  by  $Z(R)$ . A prime ideal  $P$  is said to be a *minimal prime ideal over an ideal  $I$*  if it is minimal among all prime ideals containing  $I$ . A prime ideal is said to be a *minimal prime ideal* if it is a minimal prime ideal over the zero ideal. Recall that an element  $x$  of ring  $R$  is said to be *nilpotent* if there exist positive integer  $n$  such that  $x^n = 0$ . The set of all nilpotent elements of ring  $R$  is said to be *nilradical* and it is denoted by  $nil(R)$ .

Let  $R$  be a ring. In Section 2 of this article, some results regarding diameter of  $L(\Gamma(R))$  is proved. It is proved in Theorem 2.1 that if  $\Gamma(R)$  is a complete graph then  $diam(L(\Gamma(R))) \in \{0, 1, 2\}$ . It is shown by means of an examples in Remark 2.2 that  $diam(L(\Gamma(R)))$  attains all the three values 0,1,2, when  $\Gamma(R)$  is a complete graph. In Theorem 2.3 it is proved that When  $diam(\Gamma(R)) = 2$ , then  $1 \leq diam(L(\Gamma(R))) \leq 3$ . In example 2.4, example of a ring  $R$  is given for which  $diam(\Gamma(R)) = 2$  and  $diam(L(\Gamma(R))) = 3$  and in example 2.5, example of a ring is given for which  $diam(\Gamma(R)) = 2 = diam(L(\Gamma(R)))$ .

## 2. On the diameter of $L(\Gamma(R))$

**Theorem 2.1.** *Let  $R$  be a commutative ring. If  $diam(\Gamma(R)) = 1$ , then  $diam(L(\Gamma(R))) \in \{0, 1, 2\}$ .*

*Proof.* As  $diam(\Gamma(R)) = 1$ , it follows from [2, Theorem 2.6] that  $xy = 0$  for each pair of distinct zero divisors  $x$  and  $y$  of  $R$  and  $R$  has atleast two zero divisors. So,  $R \cong \mathbb{Z}_2 \times \mathbb{Z}_2$  or  $Z(R) = P$  with  $P^2 = (0)$ .

**Case (i):**  $R \cong \mathbb{Z}_2 \times \mathbb{Z}_2$ .

Then,  $\Gamma(R)$  has only edge  $(0, 1) - (1, 0)$ . So,  $L(\Gamma(R))$  has only one vertex. So,  $diam(L(\Gamma(R))) = 0$ .

**Case (ii):**  $Z(R) = P$  with  $P^2 = (0)$ .

**Subcase (i):** If  $|P| = 3$ .

Then,  $|Z^*(R)| = |P^*| = 2$ . Let  $x, y \in Z^*(R), x \neq y$ . Then,  $\Gamma(R)$  has only one edge  $x - y$ . So,  $L(\Gamma(R))$  has only one vertex  $[x, y]$ . So,  $diam(L(\Gamma(R))) = 0$ .

**Subcase (ii):** If  $|P| = 4$ .

Then,  $|Z^*(R)| = |P^*| = 3$ . As,  $diam(\Gamma(R)) = 1$ , it follows  $\Gamma(R)$  is a triangle. So,  $L(\Gamma(R))$  is a path on two vertices. So,  $diam(L(\Gamma(R))) = 1$ .

**Subcase (iii):** If  $|P| \geq 5$ . Let  $a, b, c, d \in P^*$  and  $e_1 = [a \ b]$  and  $e_2 = [c \ d]$  be any two vertices of  $L(\Gamma(R))$ . Also, note that  $e_1$  and  $e_2$  are not adjacent in  $L(\Gamma(R))$ . So,  $diam(L(\Gamma(R))) \geq 2$ . As,  $diam(\Gamma(R)) = 1$ , it follows that  $a$  and  $c$  are adjacent in  $\Gamma(R)$ . So, we have a path  $e_1 = [a \ b] - [a \ c] - [c \ d] = e_2$  between  $e_1$  and  $e_2$ . So,  $diam(L(\Gamma(R))) \leq 2$ . Hence,  $diam(L(\Gamma(R))) = 2$ .  $\square$

**Remark 2.2.** *Here we present examples to show that in above theorem  $diam(L(\Gamma(R)))$  attains all the three number 0, 1, 2.*

*Note that the zero divisor of  $\Gamma(R)$  for  $R = \mathbb{Z}_6, \frac{F_4[x]}{(x^2)}, \mathbb{Z}_{25}$  is a complete graph. So, for  $R \in \{\mathbb{Z}_6, \frac{F_4[x]}{(x^2)}, \mathbb{Z}_{25}\}$ ,  $diam(\Gamma(R)) = 1$ . But  $diam(L(\Gamma(\mathbb{Z}_6))) = 0$ ,  $diam\left(L\left(\Gamma\left(\frac{F_4[x]}{(x^2)}\right)\right)\right) = 1$  and  $diam(L(\Gamma(\mathbb{Z}_{25}))) = 2$ .*

**Theorem 2.3.** *Let  $R$  be a commutative ring. If  $diam(\Gamma(R)) = 2$ , then  $1 \leq diam(L(\Gamma(R))) \leq 3$ .*

*Proof.* Since,  $diam(\Gamma(R)) = 2$ , it follows from [3, Theorem 2.6] that either  $R$  is reduced with exactly two minimal prime ideals and atleast three nonzero zero divisors or  $Z(R)$  is an ideal whose square is not  $(0)$  and each pair of distinct zero divisors has a nonzero annihilator.

**Case (i):**  $R$  is reduced with exactly two minimal prime ideals  $P_1$  and  $P_2$  and at least three nonzero zero divisors.

Then,  $Z(R) = P_1 \cup P_2$  and  $P_1 \cap P_2 = (0)$ .

**Subcase (i):**  $|P_1| = 2$  and  $|P_2| \geq 3$ .

Then,  $\Gamma(R)$  is a star graph  $K_{1,n}$ , where  $|P_2| = n + 1$ . So,  $L(\Gamma(R))$  is a complete graph on  $\frac{n(n+1)}{2}$ . So,  $diam(L(\Gamma(R))) = 1$ .

**Subcase(iii)**  $|P_1| \geq 3$  and  $|P_2| \geq 3$ . Then,  $\Gamma(R)$  is a complete bipartite graph with vertex partition  $Z^*(R) = V_1 \cup V_2$ , where  $V_1 = P_1 \setminus \{0\}$  and  $V_2 = P_2 \setminus \{0\}$ . Since,  $|P_1| \geq 3$  and  $|P_2| \geq 3$ , it follows that  $|V_1| \geq 2$  and  $|V_2| \geq 2$ . Let  $x_1, x \in V_1, x_1 \neq x$  and  $y_1, y \in V_2, y_1 \neq y$ . Then,  $e_1 = [x \ y], e_2 = [x_1 \ y_1]$  are vertices of  $L(\Gamma(R))$ . Note that  $e_1$  and  $e_2$  are not adjacent in  $L(\Gamma(R))$ . So,  $diam(L(\Gamma(R))) \geq 2$ . Let  $e_1 = [a \ b], e_2 = [c \ d] \in V(L(\Gamma(R)))$  and  $e_1$  and  $e_2$  are not adjacent vertices of  $V(L(\Gamma(R)))$ . As,  $a - b$  is an edge of  $\Gamma(R)$ , without loss of generality, we can assume that  $a \in V_1 = P_1 \setminus \{0\}$  and  $b \in V_2 = P_2 \setminus \{0\}$ . Similarly, we can assume that  $c \in V_1 = P_1 \setminus \{0\}$  and  $d \in V_2 = P_2 \setminus \{0\}$ . As,  $a \in V_1$  and  $d \in V_2$ , they are adjacent in  $\Gamma(R)$ . So, we have a path  $e_1 = [a \ b] - [a \ d] - [c \ d] = e_2$ . Hence,  $diam(L(\Gamma(R))) \leq 2$ . Therefore,  $diam(L(\Gamma(R))) = 2$ .

**Case (ii):**  $Z(R)$  is an ideal whose square is not  $(0)$  and each pair of zero divisors has a nonzero annihilator.

Let  $Z(R) = P, P^2 \neq (0)$ . As  $diam(\Gamma(R)) = 2$ , we can find  $x, y \in Z^*(R)$  with  $xy \neq 0$  and  $x \neq y$ . Let  $e_1 = [a \ b]$  and  $e_2 = [c \ d]$  be any two vertices of  $L(\Gamma(R))$ . Assume that  $e_1$  and  $e_2$  are not adjacent in  $L(\Gamma(R))$ . If  $ac = 0$ , then we have a path  $e_1 = [a \ b] - [a \ c] - [c \ d] = e_2$  between  $e_1$  and  $e_2$  of length of 2. Similarly,  $ac = 0$ , then we have a path  $e_1 = [a \ b] - [b \ c] - [c \ d] = e_2$  between  $e_1$  and  $e_2$  of length of 2. Similarly, in the case  $ad = 0$  and  $bd = 0$ , there is a path between  $e_1$  and  $e_2$  of length of 2. So, we can assume that  $ac \neq 0, ad \neq 0, bc \neq 0, bd \neq 0$ . Now, Since  $a$  and  $c$  are two different zero divisors, by hypothesis there exist  $y \in Z^*(R)$  such that  $ay = 0 = cy$ . Now,  $y \neq a$  as  $ac \neq 0, y \neq b$  as  $bc \neq 0, y \neq d$  as  $ad \neq 0, y \neq c$  as  $ac \neq 0$ . Hence,  $y \notin \{a, b, c, d\}$ . So, we have a path  $e_1 = [a \ b] - [a \ y] - [c \ y] - [c \ d] = e_2$  between  $e_1$  and  $e_2$  of length 3. So,  $diam(L(\Gamma(R))) \leq 3$ .  $\square$

In the following Example 2.4 we gave an example of a ring  $R$  for which  $diam(\Gamma(R)) = 2$  and  $diam(L(\Gamma(R))) = 3$ .

**Example 2.4.** Consider the ring  $R = \frac{\cup_{n=1}^{\infty} K[x_1, x_2, \dots, x_n]}{I = \langle \{x_i x_j \mid i \neq j, i, j \in \mathbb{N}\} \rangle}$ . Note that  $R$  is a reduced Ring. Let  $M = \frac{\{x_i \mid i \in \mathbb{N}\}}{I}$ . Then  $M = Z(R)$  is an ideal of  $R$  with  $M^2 = (0)$ . Let  $X_i = x_i + I$ . Note that  $e_1 = [x_1 + x_3 \ x_2 + x_4]$  and  $e_2 = [x_1 + x_2 \ x_3 + x_4]$  are vertices of  $L(\Gamma(R))$ . Note that  $e_1$  and  $e_2$  are not adjacent in  $L(\Gamma(R))$ . Now,

$$\begin{aligned} (x_1 + x_3)(x_1 + x_2) &= x_1^2 \neq 0 \\ (x_1 + x_3)(x_3 + x_4) &= x_3^2 \neq 0 \\ (x_2 + x_4)(x_1 + x_2) &= x_2^2 \neq 0 \\ (x_2 + x_4)(x_3 + x_4) &= x_4^2 \neq 0. \end{aligned}$$

So,  $diam(L(\Gamma(R))) \geq 3$ . Now, by Theorem 2.2, we have  $diam(L(\Gamma(R))) \leq 3$ . Hence,  $diam(L(\Gamma(R))) = 3$ .

In the following Example 2.5 we gave an example of a ring  $R$  for which  $diam(\Gamma(R)) = 2$  and  $diam(L(\Gamma(R))) = 2$ .

**Example 2.5.** Consider the ring  $R = \frac{K[x, y]}{(x^3)}$ , where  $K$  is a field. Then  $M = Z(R) = \frac{(x)}{(x^3)}$  is an maximal ideal of  $R$  and  $M^2 \neq (0)$ . Let  $e_1 = [a \ b]$  and  $e_2 = [c \ d]$  be any two non adjacent vertices of  $L(\Gamma(R))$ . As,  $a, b, c, d \in M$ , we have  $a = \bar{f}x$  and  $a = \bar{g}x$ . Now, since  $ab = 0$ , we have  $fg \in M$ . Hence, either  $f \in M$  or  $g \in M$ . Without loss of generality we can assume that  $f \in M$ . Hence,  $f = xs$  for some  $s \in K$ . So,  $a = x^2s$ . similarly we can assume that  $c = x^2r$  for some  $r \in K$ . So, we have  $ad = 0$ . Hence, we have a path  $e_1 = [a \ b] - [a \ d] - [c \ d] = e_2$  of length 2 between  $e_1$  and  $e_2$ . So,  $diam(L(\Gamma(R))) \leq 2$ . Now, consider the elements  $e_1 = [x \ x^2y]$  and  $e_2 = [x^2 \ xy]$  of  $L(\Gamma(R))$ . Note that  $e_1$  and  $e_2$  are not adjacent in  $L(\Gamma(R))$ . Therefore,  $diam(L(\Gamma(R))) \geq 2$ . So,  $diam(L(\Gamma(R))) = 2$ .

**Lemma 2.6.** Let  $R$  be a ring,  $Z(R)$  is an ideal of  $R$  whose square is not  $(0)$  and each pair of distinct zero divisors has a nonzero annihilator. If there exist  $a, b, c, d \in Z^*(R)$  such that  $ab = 0, cd = 0, ac \neq 0, ad \neq 0, bc \neq 0, bd \neq 0$ . Then  $d_{L(\Gamma(R))}([a \ b], [c \ d]) \geq 3$ .

*Proof.* Since,  $[a \ b]$  and  $[c \ d]$  are not adjacent in  $L(\Gamma(R))$ . So,  $d_{L(\Gamma(R))}([a \ b], [c \ d]) \geq 2$ . Suppose that there exist a path of length 2 between  $[a \ b]$  and  $[c \ d]$  in  $L(\Gamma(R))$ . Let  $e_1 = [a \ b] - [x \ y] - [c \ d]$  is a path of length 2 between  $[a \ b]$  and  $[c \ d]$  in  $L(\Gamma(R))$ . Note that  $\{x, y\} \cap \{a, b\}$  is a singleton set. Without loss of generality we can assume that  $\{x, y\} \cap \{a, b\} = \{x\}$  with  $x = a$ . Then  $\{a, y\} \cap \{c, d\} = \{y\}$ . So,  $y \in \{c, d\}$ . without loss of generality we can assume that  $y = c$ . Then  $ac = 0$ . This is in contradiction to the hypothesis. So, there is no path of length 2 between  $[a \ b]$  and  $[c \ d]$  in  $L(\Gamma(R))$ .  $d_{L(\Gamma(R))}([a \ b], [c \ d]) \geq 3$ .  $\square$

**Lemma 2.7.** *Let  $R$  be a reduced ring,  $Z(R) = P$  is an ideal of  $R$  whose square is not  $(0)$ . Let  $\{a, b, c, d\} \subseteq P^*$  and the subgraph of  $\Gamma(R)$  induced on  $\{a, b, c, d\}$  is a clique, then  $diam(L(\Gamma(R))) = 3$ .*

*Proof.* Note that  $e_1 = [a + c \ b + d]$  and  $e_2 = [a + b \ c + d]$  are vertices of  $L(\Gamma(R))$ . Also we have

$$\begin{aligned} (a + c)(a + b) &= a^2 \neq 0 \\ (a + c)(c + d) &= c^2 \neq 0 \\ (b + d)(a + b) &= b^2 \neq 0 \\ (b + d)(c + d) &= d^2 \neq 0 \end{aligned}$$

If  $a + c = b + d$ , then  $d(a + c) = d(b + d)$ . Hence,  $d^2 = 0$ . This is not possible as  $R$  is reduced. so,  $a + c \neq b + d$ . Similarly,  $a + c \neq c + d$ ,  $b + d \neq c + d$ ,  $a + b \neq c + d$ . So, from Lemma 2.6, we obtain that  $d_{L(\Gamma(R))}([a \ b], [c \ d]) \geq 3$ . Therefore,  $diam(L(\Gamma(R))) \geq 3$ . As,  $diam(\Gamma(R)) = 2$ , we have  $diam(L(\Gamma(R))) \leq 3$ . Hence,  $diam(L(\Gamma(R))) = 3$ .  $\square$

**Corollary 2.8.** *Let  $R$  be a reduced ring,  $Z(R) = P$  is an ideal of  $R$  whose square is not  $(0)$  and each pair of distinct zero divisors has a non zero annihilator. If  $\omega(\Gamma(R)) \geq 4$ , then  $diam(L(\Gamma(R))) = 3$ .*

**Lemma 2.9.** *Let  $R$  be a reduced ring with exactly three minimal prime ideals then  $diam(L(\Gamma(R))) = 2$ .*

*Proof.* Let  $P_1, P_2, P_3$  are three minimal prime ideals of  $R$ . Let  $e_1 = [a \ b]$  and  $e_2 = [a \ b]$  be any two non adjacent vertices of  $L(\Gamma(R))$ . If  $ac = 0$ , then we have a path  $e_1 = [a \ b] - [a \ c] - [c \ d]$  of length 2 between  $e_1$  and  $e_2$  in  $L(\Gamma(R))$ . Similarly, if  $ad = 0, bc = 0$  or  $bd = 0$  then we have a path of length 2 between  $e_1$  and  $e_2$  in  $L(\Gamma(R))$ . So, we assume that  $ac \neq 0$ . Without loss of generality we can assume that  $ac \notin P_1$ . Hence,  $a \notin P_1$  and  $c \notin P_1$ . Now from  $ab = 0 \in P_1$  and  $a \notin P_1$ , we have  $b \in P_1$ . Similarly from  $cd = 0$  and  $c \notin P_1$ , we have  $d \in P_1$ . Now,  $bd \neq 0$ . Hence,  $bd \notin P_2$ . So,  $b \notin P_2$  and  $d \notin P_2$ . As,  $ab = 0$  and  $cd = 0$ , we have  $a \in P_2$  and  $c \in P_2$ . Now from  $ad \neq 0$ , we obtain that  $ad \notin P_3$ . Hence,  $d \notin P_3$  and  $a \notin P_3$ . Therefore,  $c \in P_3$  and  $b \in P_3$ . So,  $c \in P_2 \cap P_3$  and  $b \in P_1 \cap P_3$ . Hence,  $bc \in P_1 \cap P_2 \cap P_3 = (0)$ . Hence,  $bc = 0$ . So, we have a path  $e_1 = [a \ b] - [b \ c] - [c \ d]$  of length 2 between  $e_1$  and  $e_2$  in  $L(\Gamma(R))$ . So,  $diam(L(\Gamma(R))) = 2$ .  $\square$

## References

- [1] D. D. Anderson and M. Naseer, *Becks coloring of a commutative ring*, J. Algebra, 159(1993), 500-514.
- [2] D.F. Anderson, R. Levy, and J. Shapiro, *Zero-divisor graphs, von Neumann regular rings, and Boolean Algebras*, J. Pure Appl. Algebra, 180(3)(2003), 221-241.
- [3] D. F. Anderson and P.S. Livingston, *The zero-divisor graph of a commutative ring*, J. Algebra, 217(1999), 434447.
- [4] M. F. Atiyah and I. G. Macdonald, *Introduction to Commutative Algebra*, Addison-Wesley Publishing Company, Reading Massachusetts, (1969).
- [5] R. Balakrishnan and K. Ranganathan, *A Textbook of Graph Theory*, Universitext Springer, (2000).

- [6] I. Beck, *Coloring of commutative rings*, J. Algebra, 116(1988), 208-226.
- [7] R. Gilmer, *Multiplicative Ideal Theory*, Marcel-Dekker, New York, (1972).
- [8] K. Nazzal and M. Ghanem, *On the Line Graph of the Zero Divisor Graph for the Ring of Gaussian Integers Modulo  $n$* , International Journal of Combinatorics, Hindawi Publication, 2012(2012), 13 pages, Article ID 957284.
- [9] Sheela Suthar and Om Prakash, *Covering of Line Graph of Zero Divisor Graph over Ring*, British Journal of Mathematics and Computer Science, 5(2015), 728-734.