



Some Common Fixed Point Results for Multivalued Mapping in Cone Metric Spaces

A.K.Dubey^{†,1}, Rohit Verma[‡] and R.P.Dubey^{*}

[†] Department of Mathematics, Bhilai Institute of Technology, Chhattisgarh, India.

[‡] Department of Mathematics, Shri Shankracharya Institute of Engineering and Technology, Durg, India.

^{*} Department of Mathematics, Dr. C.V.Raman University, Chhattisgarh, India.

Abstract : Let P be a subset of a Banach space E and P is normal and regular cone on E , we prove the existence of the common fixed point for multivalued maps in cone metric spaces. Our results generalize some well-known recent results in the literature.

Keywords : Cone metric spaces, multivalued mappings, Fixed point.

AMS Subject Classification: 47H10, 54H25.

1 Introduction

Huang and Zhang [1] generalized the notion of metric space by replacing the set of real numbers by ordered Banach space, defined a cone metric space, and obtained some fixed point theorems for contractive type mappings in a normal cone metric space. Subsequently, several other authors [2], [8], [9], [10], [11] studied the existence of common fixed point of mappings satisfying a contractive type condition in cone metric spaces. Seong Hoon Cho and Mi Sun Kim [6] have proved certain fixed point theorems by using multivalued mapping in the setting of contractive constant in metric spaces. In this paper we obtain common fixed points for a pair of multivalued mappings satisfying a generalized contractive type conditions in cone metric spaces. Recall the following definitions which are related to cone metric spaces from [1].

Definition 1.1. Let E be a real Banach space and a subset P of E is said to be a cone if it satisfies that following conditions,

(i) P is closed, non-empty, and $P \neq \{0\}$,

(ii) $ax + by \in P, a, b \geq 0$ and $x, y \in P$,

(iii) $x \in P$ and $-x \in P \Rightarrow x = 0 \Leftrightarrow P \cap (-P) = \{0\}$.

The partial ordering \leq with respect to the cone P by $x \leq y$ if and only if $y - x \in P$. If $y - x \in$ interior of P , then it is denoted by $x \ll y$. The cone P is said to be normal if a number $K > 0$ such that,

¹Corresponding author E-Mail: anilkumardby@rediffmail.com (A.K.Dubey)

for all $x, y \in E, 0 \leq x \leq y$ implies $\|x\| \leq K \|y\|$. The cone P is called regular if every increasing sequence which is bounded above is convergent and every decreasing sequence which is bounded below is convergent. The least positive number K is called the normal constant of P . There are non normal cones also. In the following, we always suppose E is a Banach space, P is a cone in E with $\text{Int } P \neq \emptyset$ and \leq is partial ordering with respect to P .

Definition 1.2. Let X be a non-empty set of E . Suppose that the map $d : X \times X \rightarrow E$ satisfies;

(i) $0 \leq d(x, y)$ for all $x, y \in X$ and $d(x, y) = 0$ if and only if $x = y$;

(ii) $d(x, y) = d(y, x)$ for all $x, y \in X$;

(iii) $d(x, y) \leq d(x, z) + d(z, y)$ for all $x, y, z \in X$.

Then d is called a cone metric on X , and (X, d) is called a cone metric space.

Example 1.3. Let $E = \mathbb{R}^2, P = \{(x, y) \in E : x, y \geq 0\} \subset \mathbb{R}^2, X = \mathbb{R}$ and $d : X \times X \rightarrow E$ defined by

$$d(x, y) = (|x - y|, \alpha|x - y|),$$

where $\alpha \geq 0$ is a constant. Then (X, d) is a cone metric space [1].

Definition 1.4. Let, (X, d) be a cone metric space, $x \in X$ and $\{x_n\}$ a sequence in X . Then

(i) $\{x_n\}$ converges to x whenever for every $c \in E$ with $0 \ll c$, there is a natural number N such that for all $d(x_n, x) \ll c$ for all $n \geq N$.

(ii) $\{x_n\}$ is a Cauchy sequence whenever for every $c \in E$ with $0 \ll c$, there is a natural number N such that $d(x_n, x_m) \ll c$ for all $n, m \geq N$.

Definition 1.5. Let (X, d) is said to be a complete cone metric space, if every Cauchy sequence is convergent in X .

Let (X, d) be a metric space. We denote by $CB(X)$ the family of non-empty closed bounded subset of X and let $C(X)$ denote the set of all non-empty compact subsets of X . Let $H(., .)$ be the Hausdorff distance on $CB(X)$. That is, for $A, B \in CB(X), H(A, B) = \max \{ \text{Sup}_{a \in A} d(a, B), \text{Sup}_{b \in B} d(A, b) \}$, where $d(a, B) = \inf \{ d(a, b) : b \in B \}$ is the distance from the point a to the subset B .

Theorem 1.6 ([7]). A multivalued mapping $T : X \rightarrow CB(X)$ is called a contraction mapping if there exists $k \in (0, 1)$ such that $H(Tx, Ty) \leq kd(x, y) \quad \forall x, y \in X$ and $x \in X$ is said to be a fixed point of T if $x \in T(x)$.

2 Fixed Point

In this section, we shall give some results which generalize [7, 10, 11].

Theorem 2.1. Let (X, d) be a complete cone metric space and let mappings $T_1, T_2 : X \rightarrow C(X)$ satisfying the following conditions;

(i) For each $x \in X, T_1(x), T_2(x) \in CB(X)$,

(ii) $H(T_1(x), T_2(y)) \leq \alpha d(x, y) + \beta [d(x, T_2(y)) + d(y, T_1(x))]$

where α, β are non negative real numbers and $\alpha + 2\beta < 1$. Then there exists $p \in X$ such that $p \in T_1(x) \cap T_2(x)$.

Proof. Let $x_0 \in X, T_1(x_0)$ is a non-empty closed bounded subset of X . Choose $x_1 \in T_1(x_0)$, for this x_1 by the same reason mentioned above $T_2(x_1)$ is non-empty closed bounded subset of X . Since $x_1 \in T_1(x_0)$ and $T_1(x_0)$ and $T_2(x_1)$ are closed bounded subset of $X, \exists x_2 \in T_2(x_1)$ such that $d(x_1, x_2) \leq H(T_1(x_0), T_2(x_1)) + q$, where $q = \max \left\{ \frac{\alpha + \beta}{1 - \beta}, \frac{\alpha + \beta}{1 - \beta} \right\}$

$$\begin{aligned} d(x_1, x_2) &\leq H(T_1(x_0), T_2(x_1)) + q \\ &\leq \alpha d(x_0, x_1) + \beta [d(x_0, T_2(x_1)) + d(x_1, T_1(x_0))] + q \\ &\leq \alpha d(x_0, x_1) + \beta [d(x_0, x_2) + d(x_1, x_1)] + q \\ &\leq \alpha d(x_0, x_1) + \beta [d(x_0, x_1) + d(x_1, x_2)] + q \\ d(x_1, x_2) &\leq \frac{\alpha + \beta}{1 - \beta} d(x_0, x_1) + q \\ d(x_1, x_2) &\leq qd(x_0, x_1) + q \end{aligned}$$

For this $x_2, T_1(x_2)$ is a non-empty closed bounded subset of X . Since $x_2 \in T_2(x_1)$ and $T_2(x_1)$ and $T_1(x_2)$ are closed bounded subset of $X, \exists x_3 \in T_1(x_2)$ such that

$$\begin{aligned} d(x_2, x_3) &\leq H(T_1(x_2), T_2(x_1)) + q^2 \\ &\leq \alpha d(x_2, x_1) + \beta [d(x_2, T_2(x_1)) + d(x_1, T_1(x_2))] + q^2 \\ &\leq \alpha d(x_1, x_2) + \beta [d(x_2, x_2) + d(x_1, x_3)] + q^2 \\ &\leq \alpha d(x_1, x_2) + \beta [d(x_1, x_2) + d(x_2, x_3)] + q^2 \\ d(x_2, x_3) &\leq \frac{\alpha + \beta}{1 - \beta} d(x_1, x_2) + q^2 \\ &\leq q \{qd(x_0, x_1) + q\} + q^2 \\ &\leq q^2 d(x_0, x_1) + 2q^2 \end{aligned}$$

Continuing this process, we get a sequence $\{x_n\}$ such that $x_{n+1} \in T_2(x_n)$ or $x_{n+1} \in T_1(x_n)$ and $d(x_{n+1}, x_n) \leq q^n d(x_0, x_1) + nq^n$.

Let $0 \ll c$ be given, choose a natural number N_1 such that $q^n d(x_0, x_1) + nq^n \ll c$ for all $n \geq N_1$ this implies $d(x_{n+1}, x_n) \ll c$.

Therefore $\{x_n\}$ is a Cauchy sequence in (X, d) is a complete cone metric space, there exists $p \in X$ such that $x_n \rightarrow p$. Choose a natural number N_2 such that

$$\begin{aligned} d(x_n, p) &\ll \frac{c(1 - \beta)}{2m(1 + \beta)} \quad \text{and} \\ d(x_{n-1}, p) &\ll \frac{c(1 - \beta)}{2m(\alpha + \beta)} \quad \text{for all } n \geq N_2. \\ d(T_1(p), p) &\leq d(p, x_n) + d(x_n, T_1(p)) \\ &\leq d(p, x_n) + H(T_2(x_{n-1}), T_1(p)) \\ &\leq d(p, x_n) + \alpha d(x_{n-1}, p) + \beta [d(x_{n-1}, T_1(p)) + d(p, T_2(x_{n-1}))] \\ &\leq d(p, x_n) + \alpha d(x_{n-1}, p) + \beta [d(x_{n-1}, T_1(p)) + d(p, x_n)] \\ &\leq d(p, x_n) + \alpha d(x_{n-1}, p) + \beta [d(x_{n-1}, p) + d(p, T_1(p)) + d(p, x_n)] \\ d(T_1(p), p) &\leq \frac{\alpha + \beta}{1 - \beta} d(x_{n-1}, p) + \frac{1 + \beta}{1 - \beta} d(x_n, p) \quad \text{for all } n \geq N_2. \end{aligned}$$

$d(T_1(p), p) \ll \frac{c}{m}$ for all $m \geq 1$, we get $\frac{c}{m} - d(T_1(p), p) \in P$, and as $m \rightarrow \infty$, we get $\frac{c}{m} \rightarrow 0$ and P is closed $-d(T_1(p), p) \in P$ but $d(T_1(p), p) \in P$. Therefore $d(T_1(p), p) = 0$ and so $p \in T_1(p)$. Similarly it can be established that $p \in T_2(p)$. Hence $p \in T_1(p) \cap T_2(p)$. \square

Corollary 2.2. *Let (X, d) be a complete cone metric space and let mappings $R, S : X \rightarrow C(X)$ satisfying the following conditions;*

- (i) For each $x \in X, R(x), S(x) \in CB(X)$,
- (ii) $H(R(x), S(y)) \leq ad(x, y) + bd(x, S(y)) + cd(y, R(x))$

where a, b, c are non-negative real numbers and $a + b + c < 1$. Then there exists $p \in X$ such that $p \in R(x) \cap S(x)$.

Proof. The symmetric property of d and the above inequality imply that

$$H(R(x), S(y)) \leq ad(x, y) + \frac{b+c}{2}[d(x, S(y)) + d(y, R(x))]$$

By substituting $R = T_1, S = T_2, a = \alpha, \frac{b+c}{2} = \beta$ in Theorem (2.1), we obtain the required result. \square

Theorem 2.3. *Let (X, d) be a complete cone metric space and let mappings $T_1, T_2 : X \rightarrow C(X)$ satisfying the following conditions;*

- (i) For each $x \in X, T_1(x), T_2(x) \in CB(X)$,
- (ii) $H(T_1(x), T_2(y)) \leq \alpha d(x, y) + \beta[d(x, T_1(x)) + d(y, T_2(y))]$

where α, β are non negative real numbers and $\alpha + 2\beta < 1$. Then there exists $p \in X$ such that $p \in T_1(x) \cap T_2(x)$.

Proof. Let $x_0 \in X, T_1(x_0)$ is a non-empty closed bounded subset of X . Choose $x_1 \in T_1(x_0)$, for this x_1 by the same reason mentioned above $T_2(x_1)$ is non-empty closed bounded subset of X .

Since $x_1 \in T_1(x_0)$ and $T_1(x_0)$ and $T_2(x_1)$ are closed bounded subset of $X, \exists x_2 \in T_2(x_1)$ such that

$$d(x_1, x_2) \leq H(T_1(x_0), T_2(x_1)) + q,$$

where $q = \max \left\{ \frac{\alpha+\beta}{1-\beta}, \frac{\alpha+\beta}{1-\beta} \right\}$

$$\begin{aligned} d(x_1, x_2) &\leq H(T_1(x_0), T_2(x_1)) + q \\ &\leq \alpha d(x_0, x_1) + \beta[d(x_0, T_1(x_0)) + d(x_1, T_2(x_1))] + q \\ &\leq \alpha d(x_0, x_1) + \beta[d(x_0, x_1) + d(x_1, x_2)] + q \\ d(x_1, x_2) &\leq \frac{\alpha + \beta}{1 - \beta} d(x_0, x_1) + q \\ d(x_1, x_2) &\leq qd(x_0, x_1) + q \end{aligned}$$

For this $x_2, T_1(x_2)$ is a non-empty closed bounded subset of X . Since $x_2 \in T_2(x_1)$ and $T_2(x_1)$ and $T_1(x_2)$ are closed bounded subset of $X, \exists x_3 \in T_1(x_2)$ such that

$$\begin{aligned} d(x_2, x_3) &\leq H(T_1(x_2), T_2(x_1)) + q^2 \\ &\leq \alpha d(x_1, x_2) + \beta[d(x_2, T_1(x_2)) + d(x_1, T_2(x_1))] + q^2 \\ &\leq \alpha d(x_1, x_2) + \beta[d(x_2, x_3) + d(x_1, x_2)] + q^2 \\ d(x_2, x_3) &\leq \frac{\alpha + \beta}{1 - \beta} d(x_1, x_2) + q^2 \\ &\leq q \{qd(x_0, x_1) + q\} + q^2 = q^2 d(x_0, x_1) + 2q^2 \end{aligned}$$

Continuing this process, we get a sequence $\{x_n\}$ such that $x_{n+1} \in T_2(x_n)$ or $x_{n+1} \in T_1(x_n)$ and $d(x_{n+1}, x_n) \leq q^n d(x_0, x_1) + nq^n$. Let $0 \ll c$ be given, choose a natural number N_1 such that $q^n d(x_0, x_1) + nq^n \ll c$ for all $n \geq N_1$ this implies $d(x_{n+1}, x_n) \ll c$. Therefore $\{x_n\}$ is a Cauchy sequence in (X, d) is a complete cone metric space, there exists $p \in X$ such that $x_n \rightarrow p$. Choose a natural number N_2 such that

$$\begin{aligned} d(x_n, p) &\ll \frac{c(1-\beta)}{2m} \text{ and} \\ d(x_{n-1}, p) &\ll \frac{c(1-\beta)}{2m \alpha} \text{ for all } n \geq N_2. \\ d(T_1(p), p) &\leq d(p, x_n) + d(x_n, T_1(p)) \\ &\leq d(p, x_n) + H(T_2(x_{n-1}), T_1(p)) \\ &\leq d(p, x_n) + \alpha d(x_{n-1}, p) + \beta[d(x_{n-1}, T_2(x_{n-1})) + d(p, T_1(p))] \\ &\leq d(p, x_n) + \alpha d(x_{n-1}, p) + \beta[d(x_{n-1}, x_n) + d(p, T_1(p))] \\ d(T_1(p), p) &\leq \frac{1}{1-\beta}d(x_n, p) + \frac{\alpha}{1-\beta}d(x_{n-1}, p) + \frac{\beta}{1-\beta}d(x_{n-1}, x_n) \text{ for all } n \geq N_2. \end{aligned}$$

$d(T_1(p), p) \ll \frac{c}{m}$ for all $m \geq 1$, we get $\frac{c}{m} - d(T_1(p), p) \in P$ and as $m \rightarrow \infty$, we get $\frac{c}{m} \rightarrow 0$ and P is closed $-d(T_1(p), p) \in P$, but $d(T_1(p), p) \in P$. Therefore $d(T_1(p), p) = 0$ and so $p \in T_1(p)$.

Similarly it can be established that $p \in T_2(p)$. Hence $p \in T_1(p) \cap T_2(p)$. \square

Corollary 2.4. *Let (X, d) be a complete cone metric space and let mappings $R, S : X \rightarrow C(X)$ satisfying the following conditions;*

- (i) For each $x \in X, R(x), S(x) \in CB(X)$,
- (ii) $H(R(x), S(y)) \leq ad(x, y) + bd(x, R(x)) + cd(y, S(y))$

where a, b, c are non-negative real numbers and $a + b + c < 1$. Then there exists $p \in X$ such that $p \in R(x) \cap S(x)$.

Proof. Proof is similar to the proof of Corollary 2.4. \square

References

- [1] L.G.Huang and X.Zhang, *Cone metric spaces and fixed point theorems of contractive mappings*, Journal of Mathematical Analysis and Applications, 332(2)(2007), 1468-1476.
- [2] Sh.Rezapour and R.Hamlbarani, *Some notes on the paper-Cone metric spaces and fixed point theorems of contractive mappings*, Journal of Mathematical Analysis and Applications, 345(2)(2008), 719-724.
- [3] H. Mohebi, *Topical functions and their properties in a class of ordered Banach spaces, in continuous optimization*, Current Trends and Modern Applications, II(2005), 343-361.
- [4] H. Mohebi, H. Sadeghi and A.M. Rubinov, *Best approximation in a class of normed spaces with star-shaped cone*, Numer. Funct. Anal. Optim., 27(3-4)(2006), 411-436.
- [5] R.K. Bose and R.N. Mukherjee, *Common fixed points of some multivalued mappings*, Tamkang J. Math., 8(2)(1977), 245-249.

- [6] Seong Hoon Cho and Mi Sun Kim, *Fixed point theorems for general contractive multivalued mappings*, J. Appl. Math. Informatics, 27(2009), 343-350.
- [7] Robert M. Brooks and Klaus Schmitt, *The contraction mapping principle and some applications*, Electronic J. of Differential equations, Monograph 09(2009).
- [8] A.K. Dubey and A. Narayan, *Cone Metric Spaces and Fixed Point Theorems for Pair of contractive maps*, Mathematica Aeterna, 2(10)(2012), 839-845.
- [9] A.K. Dubey, Rita Shukla and R.P. Dubey, *An extension of the paper Cone Metric Spaces and Fixed Point Theorems of contractive mappings*, Int. J. of Applied Mathematical Research, 2(1)(2013), 84-90.
- [10] R. Krishna Kumar and M. Marudai, *Generalization of a Fixed Point Theorem in Cone Metric Spaces*, Int. J. of Math. Analysis, 5(11)(2011), 507-512.
- [11] Ismat Beg, Akbar Azam and Muhammad Arshad, *Common Fixed Points for Maps on Topological Vector Space Valued Cone Metric Spaces*, Int. J. of Mathematics and Mathematical Sciences, (2009), Article ID 560264, 8 pages, doi:10.1155/2009/560264.