



# Solution and Ulam - Hyers Stability of an Additive - Quadratic Functional Equation in Banach Space: Hyers Direct and Fixed Point Methods

Research Article

John. M. Rassias<sup>1</sup>, M. Arunkumar<sup>2\*</sup> and P.Agilan<sup>3</sup>

1 Pedagogical Department E.E., Section of Mathematics and Informatics, National and Capodistrian University of Athens, Greece.

2 Department of Mathematics, Government Arts College, Tiruvannamalai, Tamil Nadu, India.

3 Department of Mathematics, S.K.P. Engineering College, Tiruvannamalai, TamilNadu, India.

**Abstract:** In this paper, the authors establish the general solution and generalized Ulam - Hyers stability of an additive quadratic functional equation

$$\begin{aligned} f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) \\ = 4f(x) + 8[f(y) + f(-y)] + 18[f(z) + f(-z)] \end{aligned}$$

in Banach spaces, using the Hyers direct and fixed point methods.

**MSC:** 39B52, 32B72, 32B82

**Keywords:** Additive functional equations, quadratic functional equation, mixed type functional equation, generalized Ulam - Hyers stability, fixed point.

© JS Publication.

## 1. Introduction

The investigation of stability problems for functional equations is related to the famous Ulam problem [34] (in 1940), concerning the stability of group homomorphisms, which was first solved by D. H. Hyers [14], in 1941. This stability problem was further generalized by several authors [2, 11, 29, 31, 33]. We cite also other pertinent research works [1, 3-8, 10, 13, 15, 21, 23, 28, 32]. The general solution and the generalized Hyers-Ulam stability for the **quadratic-additive type functional equation**

$$f(x + ay) + af(x - y) = f(x - ay) + af(x + y) \quad (1)$$

for any positive integer  $a$  with  $a \neq -1, 0, 1$  was discussed by K.W. Jun and H.M. Kim [18]. Also, A. Najati and M.B.Moghimi [26] investigated the generalized Hyers-Ulam-Rassias stability for the **quadratic additive functional equation** of the form

$$f(2x + y) + f(2x - y) = 2f(x + y) + 2f(x - y) + 2f(2x) - 4f(x). \quad (2)$$

\* E-mail: [annarun2002@yahoo.co.in](mailto:annarun2002@yahoo.co.in)

Infact, M.E. Gordji et. al., [12] discussed the generalized Hyers- Ulam stability of the additive - quadratic functional equation

$$f(2x + y) + f(2x - y) = f(x + y) + f(x - y) + 2f(2x) - 2f(x) \quad (3)$$

in fuzzy Banach spaces. The general solution and generalized Ulam - Hyers stability of a mixed type additive quadratic(AQ)-functional equation

$$g(x + y) + g(x - y) = 2g(x) + g(y) + g(-y) \quad (4)$$

was investigated by M. Arunkumar and J.M. Rassias [6]. Also, the general solution in vector space and generalized Ulam - Hyers stability of mixed type additive quadratic functional equation

$$f(2x \pm y \pm z) = 2f(-x \mp y \mp z) - 2f(\mp y \mp z) + f(\pm y \pm z) + 3f(x) - f(-x) \quad (5)$$

in Random Normed Space was discussed by S. Murthy et.al., [25]. Several other mixed type additive - quadratic functional equations were introduced and investigated in [9, 16, 17, 19, 20, 22, 27, 35].

In this paper, the authors establish the general solution and generalized Ulam - Hyers stability of an additive quadratic functional equation

$$\begin{aligned} f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) \\ = 4f(x) + 8[f(y) + f(-y)] + 18[f(z) + f(-z)] \end{aligned} \quad (6)$$

in Banach spaces, using the Hyers direct and fixed point methods.

In Section 2, the general solution of the functional equation (6) is given.

In Sections 3 and 4, the generalized Ulam - Hyers stability of the functional equation (6) using direct method and fixed point method is proved, respectively.

## 2. General Solution of the Functional Equation(6)

In this section, the general solution of the functional equation (6) is given. Through out this section, let us assume  $X$  and  $Y$  be vector spaces.

**Lemma 2.1.** *An odd function  $f : X \rightarrow Y$  satisfies the additive functional equation*

$$f(x + y) = f(x) + f(y) \quad (7)$$

for all  $x, y \in X$ , if and only if  $f : X \rightarrow Y$  satisfies the functional equation (6) for all  $x, y, z \in X$ .

*Proof.* Let  $f : X \rightarrow Y$  satisfy the functional equation (7). Setting  $x = y = 0$  in (7), we get  $f(0) = 0$ . Replacing  $y$  by  $x$  and  $y$  by  $2x$  in (7), we obtain

$$f(2x) = 2f(x) \quad \text{and} \quad f(3x) = 3f(x) \quad (8)$$

for all  $x \in X$ . In general for any positive integer  $a$ , we have  $f(ax) = af(x)$ .

Replacing  $y$  by  $y + z$  in (7) and using (7), we get

$$f(x + y + z) = f(x) + f(y) + f(z) \quad (9)$$

for all  $x, y, z \in X$ . Again replacing  $(x, y, z)$  by  $(x, 2y, 3z)$  in (9) and using (8), we obtain

$$f(x + 2y + 3z) = f(x) + 2f(y) + 3f(z) \tag{10}$$

for all  $x, y, z \in X$ . Setting  $y$  by  $-y$  in (10), we have

$$f(x - 2y + 3z) = f(x) + 2f(-y) + 3f(z) \tag{11}$$

for all  $x, y, z \in X$ . Again setting  $z$  by  $-z$  in (10), we get

$$f(x + 2y - 3z) = f(x) + 2f(y) + 3f(-z) \tag{12}$$

for all  $x, y, z \in X$ . Putting  $(y, z)$  by  $(-y, -z)$  in (10), we obtain

$$f(x - 2y - 3z) = f(x) + 2f(-y) + 3f(-z) \tag{13}$$

for all  $x, y, z \in X$ . Adding (10), (11), (12) and (13), we arrive

$$\begin{aligned} & f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) \\ &= 4f(x) + 4f(y) + 4f(-y) + 6f(z) + 6f(-z) \end{aligned} \tag{14}$$

for all  $x, y, z \in X$ . Adding  $4f(y) + 12f(z)$  on both sides of (14), we have

$$\begin{aligned} & f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) + 4f(y) + 12f(z) \\ &= 4f(x) + 4f(y) + 4f(-y) + 6f(z) + 6f(-z) + 4f(y) + 12f(z) \end{aligned} \tag{15}$$

for all  $x, y, z \in X$ . It follows from (15) that

$$\begin{aligned} & f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) \\ &= 4f(x) + 4f(y) + 4f(-y) + 6f(z) + 6f(-z) + 4f(y) + 12f(z) - 4f(y) - 12f(z) \end{aligned} \tag{16}$$

for all  $x, y, z \in X$ . Using oddness of  $f$  in (16), we have demonstrated our result.

Conversely, let  $f : X \rightarrow Y$  satisfy the functional equation (6). Setting  $x = y = z = 0$  in (6), we get  $f(0) = 0$ . Using oddness of  $f$  in (6), we have

$$f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) = 4f(x) \tag{17}$$

for all  $x, y, z \in X$ . Replacing  $(y, z)$  by  $(\frac{y}{2}, 0)$  in (17), we get

$$f(x + y) + f(x - y) = 2f(x) \tag{18}$$

for all  $x, y \in X$ . By Theorem 2.1 of [4], our result is demonstrated. □

**Lemma 2.2.** *An even function  $f : X \rightarrow Y$  satisfies the quadratic functional equation*

$$f(x+y) + f(x-y) = 2f(x) + 2f(y) \quad (19)$$

for all  $x, y \in X$ , if and only if  $f : X \rightarrow Y$  satisfies the functional equation (6) for all  $x, y, z \in X$ .

*Proof.* Let  $f : X \rightarrow Y$  satisfy the functional equation (19). Setting  $(x, y)$  by  $(0, 0)$  in (7), we obtain  $f(0) = 0$ . Replacing  $y$  by  $x$  and  $y$  by  $2x$  in (7), we get

$$f(2x) = 4f(x) \quad \text{and} \quad f(3x) = 9f(x) \quad (20)$$

for all  $x \in X$ . In general for any positive integer  $a$ , we have

$$f(ax) = a^2 f(x) \quad (21)$$

for all  $x \in X$ . Replacing  $y$  by  $2y + 3z$  in (19), we get

$$f(x + 2y + 3z) + f(x - 2y - 3z) = 2f(x) + 2f(2y + 3z) \quad (22)$$

for all  $x, y, z \in X$ . Again replacing  $y$  by  $-2y + 3z$  in (19), we obtain

$$f(x - 2y + 3z) + f(x + 2y - 3z) = 2f(x) + 2f(-2y + 3z) \quad (23)$$

for all  $x, y, z \in X$ . Adding (22) and (23), we arrive

$$\begin{aligned} f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) \\ = 4f(x) + 2f(2y + 3z) + 2f(-2y + 3z) \end{aligned} \quad (24)$$

for all  $x, y, z \in X$ . Using (19) in (24) and using evenness of  $f$ , we have

$$\begin{aligned} f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) \\ = 4f(x) + 2[2f(2y) + 2f(3z) - f(2y - 3z)] + 2[2f(-2y) + 2f(3z) - f(-2y - 3z)] \\ = 4f(x) + 16[f(y) + f(-y)] + 72f(z) - 2[f(2y - 3z) + f(2y + 3z)] \\ = 4f(x) + 32f(y) + 72f(z) - 2[2f(2y) + 2f(3z)] \\ = 4f(x) + 32f(y) + 72f(z) - 16f(y) - 36f(z) \\ = 4f(x) + 16f(y) + 32f(z) \\ = 4f(x) + 8[f(y) + f(-y)] + 18[f(z) + f(-z)] \end{aligned} \quad (25)$$

for all  $x, y, z \in X$ .

Conversely, assume  $f : X \rightarrow Y$  satisfies the functional equation (6). Setting  $(x, y, z)$  by  $(0, 0, 0)$  in (6), we obtain  $f(0) = 0$ .

Replacing  $z$  by 0 and using evenness of  $f$  in (6), we have

$$f(x + 2y) + f(x - 2y) = 2f(x) + 8f(y) \quad (26)$$

for all  $x, y \in X$ . Setting  $x$  by 0 in (26) and using evenness of  $f$ , we get

$$f(2y) = 4f(y) \quad (27)$$

for all  $y \in X$ . Replacing  $y$  by  $\frac{y}{2}$  in (26) and using (27), we arrive (19) as desired.  $\square$

### 3. Stability Results: Hyers Direct Method

In this section, the generalized Ulam - Hyers stability of functional equation (6), using the Hyers direct method, is provided. Now, let us consider  $X$  and  $Y$  to be a normed space and a Banach space, respectively. Define a mapping  $Df : X \rightarrow Y$  by

$$Df(x, y, z) = f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) - 4f(x) - 8[f(y) + f(-y)] - 18[f(z) + f(-z)]$$

for all  $x, y, z \in X$ .

**Theorem 3.1.** Let  $j \in \{-1, 1\}$  and  $\alpha, \beta : X^3 \rightarrow [0, \infty)$  be a function such that

$$\lim_{n \rightarrow \infty} \frac{\alpha(6^{nj}x, 6^{nj}y, 6^{nj}z)}{6^{nj}} = 0 \tag{28}$$

for all  $x, y, z \in X$ . Let  $f_a : X \rightarrow Y$  be an odd function satisfying the inequality

$$\|Df_a(x, y, z)\| \leq \alpha(x, y, z) \tag{29}$$

for all  $x, y, z \in X$ . Then there exists a unique additive mapping  $A : X \rightarrow Y$  which satisfies (6) and

$$\|f_a(x) - A(x)\| \leq \frac{1}{12} \sum_{k=\frac{1-j}{2}}^{\infty} \frac{\beta(6^{kj}x)}{6^{kj}} \tag{30}$$

where  $\beta(6^{kj}x)$  and  $A(x)$  are defined by the following two formulas

$$\beta(6^{kj}x) = 2\alpha(6^{kj}x, 6^{kj}x, 6^{kj}x) + \alpha(6^{kj}x, 0, 6^{kj}x) \tag{31}$$

$$A(x) = \lim_{n \rightarrow \infty} \frac{f_a(6^{nj}x)}{6^{nj}} \tag{32}$$

respectively, for all  $x \in X$ .

*Proof.* Assume  $j = 1$ . Replacing  $(x, y, z)$  by  $(x, x, x)$  in (29) and using oddness of  $f_a$ , we get

$$\|f_a(6x) + f_a(2x) - f_a(4x) - 4f_a(x)\| \leq \alpha(x, x, x) \tag{33}$$

for all  $x \in X$ . Again replacing  $(x, y, z)$  by  $(x, 0, x)$  in (29) and using oddness of  $f_a$ , we obtain

$$\|2f_a(4x) - 2f_a(2x) - 4f_a(x)\| \leq \alpha(x, 0, x) \tag{34}$$

for all  $x \in X$ . It follows from (33) and (34) and the triangle inequality that

$$\begin{aligned} \|2f_a(6x) - 12f_a(x)\| &\leq 2\|f_a(6x) + f_a(2x) - f_a(4x) - 4f_a(x)\| + \|2f_a(4x) - 2f_a(2x) - 4f_a(x)\| \\ &\leq 2\alpha(x, x, x) + \alpha(x, 0, x) \end{aligned} \tag{35}$$

for all  $x \in X$ . Divide the above inequality by 12, we obtain

$$\left\| \frac{f_a(6x)}{6} - f_a(x) \right\| \leq \frac{\beta(x)}{12} \tag{36}$$

where

$$\beta(x) = 2\alpha(x, x, x) + \alpha(x, 0, x)$$

for all  $x \in X$ . Now, replacing  $x$  by  $6x$  and dividing by 6 in (36), we get

$$\left\| \frac{f_a(6^2x)}{6^2} - \frac{f_a(6x)}{6} \right\| \leq \frac{\beta(6x)}{12 \cdot 6} \quad (37)$$

for all  $x \in X$ . From (36) and (37), we obtain

$$\begin{aligned} \left\| \frac{f_a(6^2x)}{6^2} - f_a(x) \right\| &\leq \left\| \frac{f_a(6x)}{6} - f_a(x) \right\| + \left\| \frac{f_a(6^2x)}{6^2} - \frac{f_a(6x)}{6} \right\| \\ &\leq \frac{1}{12} \left[ \beta(x) + \frac{\beta(6x)}{6} \right] \end{aligned} \quad (38)$$

for all  $x \in X$ . Proceeding further and using induction on a positive integer  $n$ , we get

$$\begin{aligned} \left\| \frac{f_a(6^n x)}{6^n} - f_a(x) \right\| &\leq \frac{1}{12} \sum_{k=0}^{n-1} \frac{\beta(6^k x)}{6^k} \\ &\leq \frac{1}{12} \sum_{k=0}^{\infty} \frac{\beta(6^k x)}{6^k} \end{aligned} \quad (39)$$

for all  $x \in X$ . In order to prove the convergence of the sequence

$$\left\{ \frac{f_a(6^n x)}{6^n} \right\},$$

first replace  $x$  by  $6^m x$  and then divide by  $6^m$  in (39), for any  $m, n > 0$ , and thus we deduce

$$\begin{aligned} \left\| \frac{f_a(6^{n+m} x)}{6^{n+m}} - \frac{f_a(6^m x)}{6^m} \right\| &= \frac{1}{6^m} \left\| \frac{f_a(6^n \cdot 6^m x)}{6^n} - f_a(6^m x) \right\| \\ &\leq \frac{1}{12} \sum_{k=0}^{n-1} \frac{\beta(6^{k+m} x)}{6^{k+m}} \\ &\leq \frac{1}{12} \sum_{k=0}^{\infty} \frac{\beta(6^{k+m} x)}{6^{k+m}} \\ &\rightarrow 0 \quad \text{as } m \rightarrow \infty \end{aligned}$$

for all  $x \in X$ . Hence the sequence  $\left\{ \frac{f_a(6^n x)}{6^n} \right\}$  is a Cauchy sequence. Since  $Y$  is complete, there exists a mapping  $A : X \rightarrow Y$  such that

$$A(x) = \lim_{n \rightarrow \infty} \frac{f_a(6^n x)}{6^n}, \quad \forall x \in X.$$

Letting  $n \rightarrow \infty$  in (39), we see that (30) holds for all  $x \in X$ . Claim that  $A$  satisfies (6). In fact, replacing  $(x, y, z)$  by  $(6^n x, 6^n y, 6^n z)$  and dividing by  $6^n$  in (29), we obtain

$$\frac{1}{6^n} \left\| Df_a(6^n x, 6^n y, 6^n z) \right\| \leq \frac{1}{6^n} \alpha(6^n x, 6^n y, 6^n z)$$

for all  $x, y, z \in X$ . Letting  $n \rightarrow \infty$  in the above inequality and using the definition of  $A$ , we see that

$$DA(x, y, z) = 0.$$

Hence  $A$  satisfies (6) for all  $x, y, z \in X$ . To show  $A$  is unique, let  $B$  be another additive mapping satisfying (6) and (30), then

$$\begin{aligned} \|A(x) - B(x)\| &= \frac{1}{6^n} \|A(6^n x) - B(6^n x)\| \\ &\leq \frac{1}{6^n} \{ \|A(6^n x) - f_a(6^n x)\| + \|f_a(6^n x) - B(6^n x)\| \} \\ &\leq \frac{1}{6} \sum_{k=0}^{\infty} \frac{\beta(6^{k+n} x)}{6^{(k+n)}} \\ &\rightarrow 0 \text{ as } n \rightarrow \infty \end{aligned}$$

for all  $x \in X$ . Hence  $A = B$  is unique.

For  $j = -1$ , we can prove a similar stability result. This completes the proof of the theorem. □

The following Corollary is an immediate consequence of Theorem 3.1 concerning the stability of (6).

**Corollary 3.2.** *Let  $\lambda$  and  $s$  be nonnegative real numbers. Let an odd function  $f_a : X \rightarrow Y$  satisfy the inequality*

$$\|Df_a(x, y, z)\| \leq \begin{cases} \lambda, & s < 1 \text{ or } s > 1; \\ \lambda \{ \|x\|^s + \|y\|^s + \|z\|^s \}, & 3s < 1 \text{ or } 3s > 1; \\ \lambda \|x\|^s \|y\|^s \|z\|^s, & 3s < 1 \text{ or } 3s > 1; \\ \lambda \{ \|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \} \}, & 3s < 1 \text{ or } 3s > 1; \end{cases} \quad (40)$$

for all  $x, y, z \in X$ . Then there exists a unique additive function  $A : X \rightarrow Y$  such that

$$\|f_a(x) - A(x)\| \leq \begin{cases} \frac{3\lambda}{10}, \\ \frac{4\lambda \|x\|^s}{|6 - 6^s|}, \\ \frac{\lambda \|x\|^{3s}}{|6 - 6^{3s}|}, \\ \frac{5\lambda \|x\|^{3s}}{|6 - 6^{3s}|} \end{cases} \quad (41)$$

for all  $x \in X$ .

Now we provide an example to illustrate that the functional equation (6) is not stable for  $s = 1$  in Condition (ii) of Corollary 3.2.

**Example 3.3.** *Let  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  be a function defined by*

$$\phi(x) = \begin{cases} \mu x, & \text{if } |x| < 1 \\ \mu, & \text{otherwise} \end{cases}$$

where  $\mu > 0$  is a constant, and define a function  $f_a : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f_a(x) = \sum_{n=0}^{\infty} \frac{\phi(6^n x)}{6^n} \quad \text{for all } x \in \mathbb{R}.$$

Then  $f_a$  satisfies the functional inequality

$$|Df_a(x, y, z)| \leq 72\mu (|x| + |y| + |z|) \quad (42)$$

for all  $x, y, z \in \mathbb{R}$ . Then there is no an additive mapping  $A : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\beta > 0$  such that

$$|f_a(x) - A(x)| \leq \beta|x| \quad \text{for all } x \in \mathbb{R}. \quad (43)$$

*Proof.* Now

$$|f_a(x)| \leq \sum_{n=0}^{\infty} \frac{|\phi(6^n x)|}{|6^n|} = \sum_{n=0}^{\infty} \frac{\mu}{6^n} = \frac{1}{1 - \frac{1}{6}} \mu = \frac{6\mu}{5}.$$

Therefore we see that  $f_a$  is bounded. We are now going to prove that  $f_a$  satisfies (42).

If  $x = y = z = 0$  then (42) is trivial. If  $|x| + |y| + |z| \geq 1$  then the left hand side of (42) is less than  $72\mu$ . Now suppose that  $0 < |x| + |y| + |z| < 1$ . Then there exists a positive integer  $k$  such that

$$\frac{1}{6^k} \leq |x| + |y| + |z| < \frac{1}{6^{k-1}}, \quad (44)$$

so that  $6^{k-1}|x| < 1$ ,  $6^{k-1}|y| < 1$ ,  $6^{k-1}|z| < 1$  and consequently

$$\begin{aligned} &6^{k-1}(x + 2y + 3z), 6^{k-1}(x - 2y + 3z), 6^{k-1}(x + 2y - 3z), 6^{k-1}(x - 2y - 3z), \\ &6^{k-1}(x), 6^{k-1}(-x), 6^{k-1}(y), 6^{k-1}(-y), 6^{k-1}(z), 6^{k-1}(-z) \in (-1, 1). \end{aligned}$$

Therefore for each  $n = 0, 1, \dots, k - 1$ , we have

$$\begin{aligned} &6^n(x + 2y + 3z), 6^n(x - 2y + 3z), 6^n(x + 2y - 3z), 6^n(x - 2y - 3z), \\ &6^n(x), 6^n(-x), 6^n(y), 6^n(-y), 6^n(z), 6^n(-z) \in (-1, 1). \end{aligned}$$

and

$$\begin{aligned} &\phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \\ &- 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] = 0 \end{aligned}$$

for  $n = 0, 1, \dots, k - 1$ . From the definition of  $f_a$  and (44), we obtain that

$$\begin{aligned} &|Df_a(x, y, z)| \\ &= \sum_{n=0}^{\infty} \frac{1}{6^n} \left| \phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \right. \\ &\quad \left. - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] \right| \\ &= \sum_{n=k}^{\infty} \frac{1}{6^n} \left| \phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \right. \\ &\quad \left. - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] \right| \\ &\leq \sum_{n=k}^{\infty} \frac{1}{6^n} 60\mu = 60\mu \times \frac{1}{6^k} \times \frac{6}{5} \leq 72\mu(|x| + |y| + |z|). \end{aligned}$$

Thus  $f_a$  satisfies (42) for all  $x, y, z \in \mathbb{R}$  with  $0 < |x| + |y| + |z| < 1$ .

We claim that the additive functional equation (6) is not stable for  $s = 1$  in condition (ii) of Corollary 3.2. Suppose on the contrary that there exists an additive mapping  $A : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\beta > 0$  satisfying (43). Since  $f_a$  is bounded and continuous for all  $x \in \mathbb{R}$ ,  $A$  is bounded on any open interval containing the origin and continuous at the origin. In view of Theorem 3.1,  $A$  must have the form  $A(x) = cx$  for any  $x$  in  $\mathbb{R}$ . Thus we obtain that

$$|f_a(x)| \leq (\beta + |c|)|x|. \quad (45)$$



But we can choose a positive integer  $m$  with  $m\mu > \beta + |c|$ .

If  $x \in (0, \frac{1}{6^{m-1}})$ , then  $6^n x \in (0, 1)$  for all  $n = 0, 1, \dots, m - 1$ . For this  $x$ , we get

$$f_a(x) = \sum_{n=0}^{\infty} \frac{\alpha(6^n x)}{6^n} \geq \sum_{n=0}^{m-1} \frac{\mu(6^n x)}{6^n} = m\mu x > (\beta + |c|) x$$

which contradicts (45). Therefore the additive functional equation (6) is not stable in the sense of Ulam, Hyers and Rassias if  $s = 1$ , assumed in the inequality (40). □

A counter example to illustrate the non stability in Condition (iii) of Corollary 3.2:

**Example 3.4.** Let  $s$  be such that  $0 < s < \frac{1}{3}$ . Then there is a function  $f_a : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\lambda > 0$  satisfying

$$|Df_a(x, y, z)| \leq \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{1-2s}{3}} \tag{46}$$

for all  $x, y, z \in \mathbb{R}$  and

$$\sup_{x \neq 0} \frac{|f_a(x) - A(x)|}{|x|} = +\infty \tag{47}$$

for every additive mapping  $A : \mathbb{R} \rightarrow \mathbb{R}$ .

*Proof.* If we take

$$f_a(x) = \begin{cases} x \ln |x|, & \text{if } x \neq 0, \\ 0, & \text{if } x = 0. \end{cases}$$

Then from the relation (47), it follows that

$$\begin{aligned} \sup_{x \neq 0} \frac{|f_a(x) - A(x)|}{|x|} &\geq \sup_{\substack{n \in \mathbb{N} \\ n \neq 0}} \frac{|f_a(n) - A(n)|}{|n|} \\ &= \sup_{\substack{n \in \mathbb{N} \\ n \neq 0}} \frac{|n \ln |n| - n A(1)|}{|n|} \\ &= \sup_{\substack{n \in \mathbb{N} \\ n \neq 0}} |\ln |n| - A(1)| = \infty. \end{aligned}$$

We have to prove (46) is true.

**Case (i):** If  $x, y, z > 0$  than  $x + 2z + 3z > 0, x - 2z + 3z > 0,$

$x + 2z - 3z > 0, x - 2z - 3z > 0$  and therefore (46) becomes,

$$\begin{aligned} &|f_a(x + 2y + 3z) + f_a(x - 2y + 3z) + f_a(x + 2y - 3z) + f_a(x - 2y - 3z) \\ &\quad - 4f_a(x) - 8[f_a(y) + f_a(-y)] - 18[f_a(z) + f_a(-z)]|. \\ &= |(x + 2y + 3z) \ln |x + 2y + 3z| + (x - 2y + 3z) \ln |x - 2y + 3z| + (x + 2y - 3z) \ln |x + 2y - 3z| \\ &\quad + (x - 2y - 3z) \ln |x - 2y - 3z| - 4x \ln |x| - 8[y \ln |y| - y \ln |-y|] - 18[z \ln |z| - z \ln |-z|]|. \end{aligned}$$

Set  $x = u, y = v, z = w$  it follows that

$$\begin{aligned}
 & |f_a(x + 2y + 3z) + f_a(x - 2y + 3z) + f_a(x + 2y - 3z) + f_a(x - 2y - 3z) \\
 & \quad - 4f_a(x) - 8[f_a(y) + f_a(-y)] - 18[f_a(z) + f_a(-z)]| \\
 & = |(x + 2y + 3z) \ln |x + 2y + 3z| + (x - 2y + 3z) \ln |x - 2y + 3z| + (x + 2y - 3z) \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z) \ln |x - 2y - 3z| - 4x \ln |x| - 8[y \ln |y| - y \ln |-y|] - 18[z \ln |z| - z \ln |-z|]|. \\
 & = |(u + 2v + 3w) \ln |u + 2v + 3w| + (u - 2v + 3w) \ln |u - 2v + 3w| + (u + 2v - 3w) \ln |u + 2v - 3w| \\
 & \quad + (u - 2v - 3w) \ln |u - 2v - 3w| - 4u \ln |u| - 8[v \ln |v| - v \ln |-v|] - 18[w \ln |w| - w \ln |-w|]|. \\
 & |f_a(u + 2v + 3w) + f_a(u - 2v + 3w) + f_a(u + 2v - 3w) + f_a(u - 2v - 3w) \\
 & \quad - 4f_a(u) - 8[f_a(v) + f_a(-v)] - 18[f_a(w) + f_a(-w)]|. \\
 & \leq \lambda |u|^{\frac{s}{3}} |v|^{\frac{s}{3}} |w|^{\frac{1-2s}{3}} = \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{1-2s}{3}}.
 \end{aligned}$$

**Case (ii):** If  $x, y, z < 0$  than  $x + 2z + 3z > 0, x - 2z + 3z > 0,$

$x + 2z - 3z > 0, x - 2z - 3z > 0$  and therefore (46) becomes,

$$\begin{aligned}
 & |f_a(x + 2y + 3z) + f_a(x - 2y + 3z) + f_a(x + 2y - 3z) + f_a(x - 2y - 3z) \\
 & \quad - 4f_a(x) - 8[f_a(y) + f_a(-y)] - 18[f_a(z) + f_a(-z)]|. \\
 & = |(x + 2y + 3z) \ln |x + 2y + 3z| + (x - 2y + 3z) \ln |x - 2y + 3z| + (x + 2y - 3z) \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z) \ln |x - 2y - 3z| - 4x \ln |x| - 8[y \ln |y| - y \ln |-y|] - 18[z \ln |z| - z \ln |-z|]|.
 \end{aligned}$$

Set  $x = -u, y = -v, z = -w$  it follows that

$$\begin{aligned}
 & |f_a(x + 2y + 3z) + f_a(x - 2y + 3z) + f_a(x + 2y - 3z) + f_a(x - 2y - 3z) \\
 & \quad - 4f_a(x) - 8[f_a(y) + f_a(-y)] - 18[f_a(z) + f_a(-z)]| \\
 & = |(x + 2y + 3z) \ln |x + 2y + 3z| + (x - 2y + 3z) \ln |x - 2y + 3z| + (x + 2y - 3z) \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z) \ln |x - 2y - 3z| - 4x \ln |x| - 8[y \ln |y| - y \ln |-y|] - 18[z \ln |z| - z \ln |-z|]|. \\
 & = |(-u - 2v - 3w) \ln |-u - 2v - 3w| + (-u + 2v - 3w) \ln |-u + 2v - 3w| \\
 & \quad + (-u - 2v + 3w) \ln |-u - 2v + 3w| + (-u + 2v + 3w) \ln |-u + 2v + 3w| \\
 & \quad + 4u \ln |-u| - 8[-v \ln |-v| + v \ln |v|] - 18[-w \ln |-w| + w \ln |w|]|. \\
 & |f_a(-u - 2v - 3w) + f_a(-u + 2v - 3w) + f_a(-u - 2v + 3w) + f_a(-u + 2v + 3w) \\
 & \quad - 4f_a(-u) - 8[f_a(-v) + f_a(v)] - 18[f_a(-w) + f_a(w)]|. \\
 & \leq \lambda |-u|^{\frac{s}{3}} |-v|^{\frac{s}{3}} |-w|^{\frac{1-2s}{3}} = \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{1-2s}{3}}.
 \end{aligned}$$

**Case (iii):** If  $x > 0, y < 0, z < 0$  than  $x + 2z + 3z < 0, x - 2z + 3z < 0,$

$x + 2z - 3z < 0, x - 2z - 3z < 0$  and therefore (46) becomes,

$$\begin{aligned}
 & |f_a(x + 2y + 3z) + f_a(x - 2y + 3z) + f_a(x + 2y - 3z) + f_a(x - 2y - 3z) \\
 & \quad - 4f_a(x) - 8[f_a(y) + f_a(-y)] - 18[f_a(z) + f_a(-z)]|. \\
 & = |(x + 2y + 3z) \ln |x + 2y + 3z| + (x - 2y + 3z) \ln |x - 2y + 3z| + (x + 2y - 3z) \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z) \ln |x - 2y - 3z| - 4x \ln |x| - 8[y \ln |y| - y \ln |-y|] - 18[z \ln |z| - z \ln |-z|]|.
 \end{aligned}$$

Set  $x = u, y = -v, z = -w$  it follows that

$$\begin{aligned}
 & |f_a(x + 2y + 3z) + f_a(x - 2y + 3z) + f_a(x + 2y - 3z) + f_a(x - 2y - 3z) \\
 & \quad - 4f_a(x) - 8[f_a(y) + f_a(-y)] - 18[f_a(z) + f_a(-z)]| \\
 & = |(x + 2y + 3z) \ln |x + 2y + 3z| + (x - 2y + 3z) \ln |x - 2y + 3z| + (x + 2y - 3z) \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z) \ln |x - 2y - 3z| - 4x \ln |x| - 8[y \ln |y| - y \ln |-y|] - 18[z \ln |z| - z \ln |-z|]| \\
 & = |(u - 2v - 3w) \ln |u - 2v - 3w| + (u + 2v - 3w) \ln |u + 2v - 3w| + (u - 2v + 3w) \ln |u - 2v + 3w| \\
 & \quad + (u + 2v + 3w) \ln |u + 2v + 3w| - 4u \ln |u| - 8[-v \ln |-v| + v \ln |v|] - 18[-w \ln |-w| + w \ln |w|]| \\
 & |f_a(u - 2v - 3w) + f_a(u + 2v - 3w) + f_a(u - 2v + 3w) + f_a(u + 2v + 3w) \\
 & \quad - 4f_a(u) - 8[f_a(-v) + f_a(v)] - 18[f_a(-w) + f_a(w)]| \\
 & \leq \lambda |u|^{\frac{s}{3}} |-v|^{\frac{s}{3}} |-w|^{\frac{1-2s}{3}} = \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{1-2s}{3}}.
 \end{aligned}$$

**Case (iv):** If  $x < 0, y > 0, z > 0$  than  $x + 2z + 3z < 0, x - 2z + 3z < 0,$   
 $x + 2z - 3z < 0, x - 2z - 3z < 0$  and therefore (46) becomes,

$$\begin{aligned}
 & |f_a(x + 2y + 3z) + f_a(x - 2y + 3z) + f_a(x + 2y - 3z) + f_a(x - 2y - 3z) \\
 & \quad - 4f_a(x) - 8[f_a(y) + f_a(-y)] - 18[f_a(z) + f_a(-z)]| \\
 & = |(x + 2y + 3z) \ln |x + 2y + 3z| + (x - 2y + 3z) \ln |x - 2y + 3z| + (x + 2y - 3z) \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z) \ln |x - 2y - 3z| - 4x \ln |x| - 8[y \ln |y| - y \ln |-y|] - 18[z \ln |z| - z \ln |-z|]|.
 \end{aligned}$$

Set  $x = -u, y = v, z = w$  it follows that

$$\begin{aligned}
 & |f_a(x + 2y + 3z) + f_a(x - 2y + 3z) + f_a(x + 2y - 3z) + f_a(x - 2y - 3z) \\
 & \quad - 4f_a(x) - 8[f_a(y) + f_a(-y)] - 18[f_a(z) + f_a(-z)]| \\
 & = |(x + 2y + 3z) \ln |x + 2y + 3z| + (x - 2y + 3z) \ln |x - 2y + 3z| + (x + 2y - 3z) \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z) \ln |x - 2y - 3z| - 4x \ln |x| - 8[y \ln |y| - y \ln |-y|] - 18[z \ln |z| - z \ln |-z|]| \\
 & = |(-u + 2v + 3w) \ln |-u + 2v + 3w| + (-u - 2v + 3w) \ln |-u - 2v + 3w| \\
 & \quad + (-u + 2v - 3w) \ln |-u + 2v - 3w| + (-u - 2v - 3w) \ln |-u - 2v - 3w| \\
 & \quad + 4u \ln |u| - 8[v \ln |v| - v \ln |-v|] - 18[w \ln |w| - w \ln |-w|]| \\
 & |f_a(-u + 2v + 3w) + f_a(-u - 2v + 3w) + f_a(-u + 2v - 3w) + f_a(-u - 2v - 3w) \\
 & \quad - 4f_a(-u) - 8[f_a(v) + f_a(-v)] - 18[f_a(w) + f_a(-w)]| \\
 & \leq \lambda |-u|^{\frac{s}{3}} |v|^{\frac{s}{3}} |w|^{\frac{1-2s}{3}} = \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{1-2s}{3}}.
 \end{aligned}$$

**Case (v):** If  $x = y = z = 0$  in (46) then it is trivial. □

Now we will provide an example to illustrate that the functional equation (6) is not stable for  $s = \frac{1}{3}$  in Condition (iv) of Corollary 3.2.

**Example 3.5.** Let  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  be a function defined by

$$\phi(x) = \begin{cases} \mu x, & \text{if } |x| < \frac{1}{3} \\ \frac{\mu}{3}, & \text{otherwise} \end{cases}$$

where  $\mu > 0$  is a constant, and define a function  $f_a : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f_a(x) = \sum_{n=0}^{\infty} \frac{\phi(6^n x)}{6^n} \quad \text{for all } x \in \mathbb{R}.$$

Then  $f_a$  satisfies the functional inequality

$$|Df_a(x, y, z)| \leq 24\mu \left( |x|^{\frac{1}{3}} |y|^{\frac{1}{3}} |z|^{\frac{1}{3}} + |x| + |y| + |z| \right), \quad (48)$$

for all  $x, y, z \in \mathbb{R}$ . Then there is no an additive mapping  $A : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\beta > 0$  such that

$$|f_a(x) - A(x)| \leq \beta|x| \quad \text{for all } x \in \mathbb{R}. \quad (49)$$

*Proof.* Now

$$|f_a(x)| \leq \sum_{n=0}^{\infty} \frac{|\phi(6^n x)|}{|6^n|} = \sum_{n=0}^{\infty} \frac{\mu}{3} \times \frac{1}{6^n} = \frac{2\mu}{5}.$$

Therefore we see that  $f_a$  is bounded. We are now going to prove that  $f_a$  satisfies (48).

If  $x = y = z = 0$  then (48) is trivial. If  $|x|^{\frac{1}{3}} |y|^{\frac{1}{3}} |z|^{\frac{1}{3}} + |x| + |y| + |z| \geq \frac{1}{6}$ , then the left hand side of (48) is less than  $24\mu$ .

Now suppose that  $0 < |x|^{\frac{1}{3}} |y|^{\frac{1}{3}} |z|^{\frac{1}{3}} + |x| + |y| + |z| < \frac{1}{6}$ . Then there exists a positive integer  $k$  such that

$$\frac{1}{6^k} \leq |x|^{\frac{1}{3}} |y|^{\frac{1}{3}} |z|^{\frac{1}{3}} + |x| + |y| + |z| < \frac{1}{6^{k+1}}, \quad (50)$$

so that  $6^{k-1}|x|^{\frac{1}{3}} |y|^{\frac{1}{3}} |z|^{\frac{1}{3}} < \frac{1}{6}$ ,  $6^{k-1}|x| < \frac{1}{6}$ ,  $6^{k-1}|y| < \frac{1}{6}$ ,  $6^{k-1}|z| < \frac{1}{6}$  and consequently

$$\begin{aligned} &6^{k-1}(x + 2y + 3z), 6^{k-1}(x - 2y + 3z), 6^{k-1}(x + 2y - 3z), 6^{k-1}(x - 2y - 3z), \\ &6^{k-1}(x), 6^{k-1}(-x), 6^{k-1}(y), 6^{k-1}(-y), 6^{k-1}(z), 6^{k-1}(-z) \in \left( \frac{-1}{6}, \frac{1}{6} \right). \end{aligned}$$

Therefore for each  $n = 0, 1, \dots, k-1$ , we have

$$\begin{aligned} &6^n(x + 2y + 3z), 6^n(x - 2y + 3z), 6^n(x + 2y - 3z), 6^n(x - 2y - 3z), \\ &6^n(x), 6^n(-x), 6^n(y), 6^n(-y), 6^n(z), 6^n(-z) \in \left( \frac{-1}{6}, \frac{1}{6} \right) \end{aligned}$$

and

$$\begin{aligned} &\phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \\ &- 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] = 0 \end{aligned}$$

for  $n = 0, 1, \dots, k-1$ . From the definition of  $f_a$  and (50), we obtain that

$$\begin{aligned} &\left| Df_a(x, y, z) \right| \\ &= \sum_{n=0}^{\infty} \frac{1}{6^n} \left| \phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \right. \\ &\quad \left. - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] \right| \\ &= \sum_{n=k}^{\infty} \frac{1}{6^n} \left| \phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \right. \\ &\quad \left. - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] \right| \\ &\leq \sum_{n=k}^{\infty} \frac{1}{6^n} \frac{60\mu}{3} = \frac{60\mu}{3} \times \frac{1}{6^k} \times \frac{6}{5} \leq 24\mu \left( |x|^{\frac{1}{3}} |y|^{\frac{1}{3}} |z|^{\frac{1}{3}} + |x| + |y| + |z| \right). \end{aligned}$$

Thus  $f_a$  satisfies (48) for all  $x, y, z \in \mathbb{R}$  with  $0 < |x|^{\frac{1}{3}}|y|^{\frac{1}{3}}|z|^{\frac{1}{3}} + |x| + |y| + |z| < \frac{1}{6}$ .

We claim that the additive functional equation (6) is not stable for  $s = \frac{1}{3}$  in condition (iv) of Corollary 3.2. Suppose on the contrary that there exists an additive mapping  $A : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\beta > 0$  satisfying (49). Since  $f_a$  is bounded and continuous for all  $x \in \mathbb{R}$ ,  $A$  is bounded on any open interval containing the origin and continuous at the origin. In view of Theorem 3.1,  $A$  must have the form  $A(x) = cx$  for any  $x$  in  $\mathbb{R}$ . Thus we obtain that

$$|f_a(x)| \leq (\beta + |c|)|x|. \tag{51}$$

But we can choose a positive integer  $m$  with  $m\mu > \beta + |c|$ .

If  $x \in (0, \frac{1}{6^{m-1}})$ , then  $6^n x \in (0, \frac{1}{6})$  for all  $n = 0, 1, \dots, m-1$ . For this  $x$ , we get

$$f_a(x) = \sum_{n=0}^{\infty} \frac{\alpha(6^n x)}{6^n} \geq \sum_{n=0}^{m-1} \frac{\mu(6^n x)}{6^n} = m\mu x > (\beta + |c|)x$$

which contradicts (51). Therefore the additive functional equation (6) is not stable in the sense of Ulam, Hyers and Rassias if  $s = \frac{1}{3}$ , assumed in the inequality (40). □

**Theorem 3.6.** Let  $j \in \{-1, 1\}$  and  $\alpha, \beta : X^3 \rightarrow [0, \infty)$  be a function such that

$$\lim_{n \rightarrow \infty} \frac{\alpha(6^{nj}x, 6^{nj}y, 6^{nj}z)}{36^{nj}} = 0 \tag{52}$$

for all  $x, y, z \in X$ . Let  $f_q : X \rightarrow Y$  be an even function satisfying the inequality

$$\|Df_q(x, y, z)\| \leq \alpha(x, y, z) \tag{53}$$

for all  $x, y, z \in X$ . Then there exists a unique quadratic mapping  $Q : X \rightarrow Y$  which satisfies (6) and

$$\|f_q(x) - Q(x)\| \leq \frac{1}{72} \sum_{k=\frac{1-j}{2}}^{\infty} \frac{\beta(6^{kj}x)}{36^{kj}} \tag{54}$$

where  $\beta(6^{kj}x)$  and  $Q(x)$  are defined by the two relations:

$$\beta(6^{kj}x) = 2\alpha(6^{kj}x, 6^{kj}x, 6^{kj}x) + \alpha(6^{kj}x, 0, 6^{kj}x) \tag{55}$$

$$Q(x) = \lim_{n \rightarrow \infty} \frac{f_q(6^{nj}x)}{36^{nj}} \tag{56}$$

respectively, for all  $x \in X$ .

*Proof.* Assume  $j = 1$ . Replacing  $(x, y, z)$  by  $(x, x, x)$  in (53) and using evenness of  $f_q$ , we get

$$\|f_q(6x) + f_q(2x) + f_q(4x) - 56f_q(x)\| \leq \alpha(x, x, x) \tag{57}$$

for all  $x \in X$ . Again replacing  $(x, y, z)$  by  $(x, 0, x)$  in (53) and using evenness of  $f_q$ , we obtain

$$\|2f_q(4x) + 2f_q(2x) - 40f_q(x)\| \leq \alpha(x, 0, x) \tag{58}$$

for all  $x \in X$ . It follows from (57) and (58) that

$$\begin{aligned} \|2f_q(6x) - 72f_q(x)\| &\leq 2\|f_q(6x) + f_q(2x) + f_q(4x) - 56f_q(x)\| \\ &\quad + \|2f_q(4x) + 2f_q(2x) - 40f_q(x)\| \\ &\leq 2\alpha(x, x, x) + \alpha(x, 0, x) \end{aligned} \quad (59)$$

for all  $x \in X$ . Dividing the above inequality by 72, we arrive

$$\left\| \frac{f_q(6x)}{36} - f_q(x) \right\| \leq \frac{\beta(x)}{72} \quad (60)$$

where

$$\beta(x) = 2\alpha(x, x, x) + \alpha(x, 0, x)$$

for all  $x \in X$ . The rest of the proof is similar to that one of Theorem 3.1.  $\square$

The following Corollary is an immediate consequence of Theorem 3.6 concerning the stability of (6).

**Corollary 3.7.** *Let  $\lambda$  and  $s$  be nonnegative real numbers. Let an even function  $f_q : X \rightarrow Y$  satisfy the inequality*

$$\|Df_q(x, y, z)\| \leq \begin{cases} \lambda, & s < 2 \quad \text{or} \quad s > 2; \\ \lambda\{\|x\|^s + \|y\|^s + \|z\|^s\}, & 3s < 2 \quad \text{or} \quad 3s > 2; \\ \lambda\|x\|^s\|y\|^s\|z\|^s, & 3s < 2 \quad \text{or} \quad 3s > 2; \\ \lambda\{\|x\|^s\|y\|^s\|z\|^s + \{\|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s}\}\}, & 3s < 2 \quad \text{or} \quad 3s > 2; \end{cases} \quad (61)$$

for all  $x, y, z \in X$ . Then there exists a unique quadratic function  $Q : X \rightarrow Y$  such that

$$\|f_q(x) - Q(x)\| \leq \begin{cases} \frac{3\lambda}{70}, \\ \frac{4\lambda\|x\|^s}{|36 - 6^s|}, \\ \frac{\lambda\|x\|^{3s}}{|36 - 6^{3s}|}, \\ \frac{5\lambda\|x\|^{3s}}{|36 - 6^{3s}|} \end{cases} \quad (62)$$

for all  $x \in X$ .

Now we provide an example to illustrate that the functional equation (6) is not stable for  $s = 2$  in Condition (ii) of Corollary 3.7.

**Example 3.8.** *Let  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  be a function defined by*

$$\alpha(x) = \begin{cases} \mu x^2, & \text{if } |x| < 2 \\ \mu, & \text{otherwise} \end{cases}$$

where  $\mu > 0$  is a constant, and define a function  $f_q : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f_q(x) = \sum_{n=0}^{\infty} \frac{\phi(6^n x)}{36^n} \quad \text{for all } x \in \mathbb{R}.$$

Then  $f_q$  satisfies the functional inequality

$$|Df_q(x, y, z)| \leq \left( \frac{12\mu \times 36^2}{7} \right) (|x|^2 + |y|^2 + |z|^2) \quad (63)$$

for all  $x, y, z \in \mathbb{R}$ . Then there is no a quadratic mapping  $Q : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\beta > 0$  such that

$$|f_q(x) - Q(x)| \leq \beta|x|^2 \quad \text{for all } x \in \mathbb{R}. \quad (64)$$

*Proof.* Now

$$|f_q(x)| \leq \sum_{n=0}^{\infty} \frac{|\phi(6^n x)|}{|36^n|} = \sum_{n=0}^{\infty} \frac{\mu}{36^n} = \frac{36\mu}{35}.$$

Therefore we see that  $f_q$  is bounded. We are going to prove that  $f_q$  satisfies (63).

If  $x = y = z = 0$  then (63) is trivial. If  $|x|^2 + |y|^2 + |z|^2 \geq \frac{1}{36}$  then the left hand side of (63) is less than  $(\frac{12\mu \times 36}{7})$ . Now suppose that  $0 < |x|^2 + |y|^2 + |z|^2 < \frac{1}{36}$ . Then there exists a positive integer  $k$  such that

$$\frac{1}{36^{k+2}} \leq |x|^2 + |y|^2 + |z|^2 < \frac{1}{36^{k+1}}, \quad (65)$$

so that  $6^{k-1}|x|^2 < \frac{1}{36}$ ,  $6^{k-1}|y|^2 < \frac{1}{36}$ ,  $6^{k-1}|z|^2 < \frac{1}{36}$  and consequently

$$6^{k-1}(x + 2y + 3z), 6^{k-1}(x - 2y + 3z), 6^{k-1}(x + 2y - 3z), 6^{k-1}(x - 2y - 3z), \\ 6^{k-1}(x), 6^{k-1}(-x), 6^{k-1}(y), 6^{k-1}(-y), 6^{k-1}(z), 6^{k-1}(-z) \in \left( -\frac{1}{6}, \frac{1}{6} \right).$$

Therefore for each  $n = 0, 1, \dots, k - 1$ , we have

$$6^n(x + 2y + 3z), 6^n(x - 2y + 3z), 6^n(x + 2y - 3z), 6^n(x - 2y - 3z), \\ 6^n(x), 6^n(-x), 6^n(y), 6^n(-y), 6^n(z), 6^n(-z) \in \left( -\frac{1}{6}, \frac{1}{6} \right).$$

and

$$\phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \\ - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] = 0$$

for  $n = 0, 1, \dots, k - 1$ . From the definition of  $f_q$  and (65), we obtain that

$$\begin{aligned} & \left| Df_q(x, y, z) \right| \\ &= \sum_{n=0}^{\infty} \frac{1}{36^n} \left| \phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \right. \\ & \quad \left. - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] \right| \\ &= \sum_{n=k}^{\infty} \frac{1}{36^n} \left| \phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \right. \\ & \quad \left. - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] \right| \\ &\leq \sum_{n=k}^{\infty} \frac{1}{6^n} 60\mu = 60 \mu \times \frac{1}{36^k} \times \frac{36}{35} \leq \left( \frac{12\mu \times 36^2}{7} \right) (|x|^2 + |y|^2 + |z|^2). \end{aligned}$$

Thus  $f_q$  satisfies (63) for all  $x, y, z \in \mathbb{R}$  with  $0 < |x|^2 + |y|^2 + |z|^2 < \frac{1}{6}$ .

We claim that the additive functional equation (6) is not stable for  $s = 2$  in condition (ii) of Corollary 3.2. Suppose on the contrary that there exists an additive mapping  $Q : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\beta > 0$  satisfying (64). Since  $f_q$  is bounded and continuous for all  $x \in \mathbb{R}$ ,  $Q$  is bounded on any open interval containing the origin and continuous at the origin. In view of Theorem 3.1,  $Q$  must have the form  $Q(x) = cx^2$  for any  $x$  in  $\mathbb{R}$ . Thus we obtain that

$$|f_q(x)| \leq (\beta + |c|) |x|^2. \quad (66)$$

But we can choose a positive integer  $m$  with  $m\mu > \beta + |c|$ .

If  $x \in (0, \frac{1}{6^{m-1}})$ , then  $6^n x \in (0, \frac{1}{6})$  for all  $n = 0, 1, \dots, m-1$ . For this  $x$ , we get

$$f_q(x) = \sum_{n=0}^{\infty} \frac{\alpha(6^n x)}{36^n} \geq \sum_{n=0}^{m-1} \frac{\mu(6^n x)^2}{36^n} = m\mu x^2 > (\beta + |c|) x^2,$$

which contradicts (66). Therefore the additive functional equation (6) is not stable in the sense of Ulam, Hyers and Rassias if  $s = 2$ , assumed in the inequality (40).  $\square$

A counter example to illustrate the non stability in Condition (iii) of Corollary 3.7:

**Example 3.9.** Let  $s$  be such that  $0 < s < \frac{2}{3}$ . Then there is a function  $f_q : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\lambda > 0$  satisfying

$$|Df_q(x, y, z)| \leq \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{2-2s}{3}} \quad (67)$$

for all  $x, y, z \in \mathbb{R}$  and

$$\sup_{x \neq 0} \frac{|f_q(x) - Q(x)|}{|x|^2} = +\infty \quad (68)$$

for every quadratic mapping  $Q : \mathbb{R} \rightarrow \mathbb{R}$ .

*Proof.* If we take

$$f_q(x) = \begin{cases} x^2 \ln |x|, & \text{if } x \neq 0, \\ 0, & \text{if } x = 0, \end{cases}$$

then from the relation (68), it follows that

$$\begin{aligned} \sup_{x \neq 0} \frac{|f_q(x) - Q(x)|}{|x|^2} &\geq \sup_{\substack{n \in \mathbb{N} \\ n \neq 0}} \frac{|f(n) - Q(n)|}{|n|^2} \\ &= \sup_{\substack{n \in \mathbb{N} \\ n \neq 0}} \frac{|n^2(1)^2 \ln |n| - n^2 Q(1)|}{|n|^2} \\ &= \sup_{\substack{n \in \mathbb{N} \\ n \neq 0}} |\ln |n| - Q(1)| = \infty. \end{aligned}$$

We have to prove that (67) is true.

**Case (i):** If  $x, y, z > 0$  than  $x + 2z + 3z > 0, x - 2z + 3z > 0,$

$x + 2z - 3z > 0, x - 2z - 3z > 0$  and therefore (46) becomes,

$$\begin{aligned} &|f_q(x + 2y + 3z) + f_q(x - 2y + 3z) + f_q(x + 2y - 3z) + f_q(x - 2y - 3z) \\ &\quad - 4f_q(x) - 8[f_q(y) + f_q(-y)] - 18[f_q(z) + f_q(-z)]|. \\ &= |(x + 2y + 3z)^2 \ln |x + 2y + 3z| + (x - 2y + 3z)^2 \ln |x - 2y + 3z| + (x + 2y - 3z)^2 \ln |x + 2y - 3z| \\ &\quad + (x - 2y - 3z)^2 \ln |x - 2y - 3z| - 4x^2 \ln |x| - 8[y^2 \ln |y| + y^2 \ln |-y|] - 18[z^2 \ln |z| + z^2 \ln |-z|]|. \end{aligned}$$



Set  $x = u, y = v, z = w$  it follows that

$$\begin{aligned}
 & |f_q(x + 2y + 3z) + f_q(x - 2y + 3z) + f_q(x + 2y - 3z) + f_q(x - 2y - 3z) \\
 & \quad - 4f_q(x) - 8[f_q(y) + f_q(-y)] - 18[f_q(z) + f_q(-z)]| \\
 & = |(u + 2v + 3w)^2 \ln |u + 2v + 3w| + (u - 2v + 3w)^2 \ln |u - 2v + 3w| + (u + 2v - 3w)^2 \ln |u + 2v - 3w| \\
 & \quad + (u - 2v - 3w)^2 \ln |u - 2v - 3w| - 4u^2 \ln |u| - 8[v^2 \ln |v| + v^2 \ln |-v|] - 18[w^2 \ln |w| + w^2 \ln |-w|]|. \\
 & |f_q(u + 2v + 3w) + f_q(u - 2v + 3w) + f_q(u + 2v - 3w) + f_q(u - 2v - 3w) \\
 & \quad - 4f_q(u) - 8[f_q(v) + f_q(-v)] - 18[f_q(w) + f_q(-w)]|. \\
 & \leq \lambda |u|^{\frac{s}{3}} |v|^{\frac{s}{3}} |w|^{\frac{2-2s}{3}} = \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{2-2s}{3}}.
 \end{aligned}$$

**Case (ii):** If  $x, y, z < 0$  than  $x + 2z + 3z > 0, x - 2z + 3z > 0,$

$x + 2z - 3z > 0, x - 2z - 3z > 0$  and therefore (46) becomes,

$$\begin{aligned}
 & |f_q(x + 2y + 3z) + f_q(x - 2y + 3z) + f_q(x + 2y - 3z) + f_q(x - 2y - 3z) \\
 & \quad - 4f_q(x) - 8[f_q(y) + f_q(-y)] - 18[f_q(z) + f_q(-z)]|. \\
 & = |(x + 2y + 3z)^2 \ln |x + 2y + 3z| + (x - 2y + 3z)^2 \ln |x - 2y + 3z| + (x + 2y - 3z)^2 \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z)^2 \ln |x - 2y - 3z| - 4x^2 \ln |x| - 8[y^2 \ln |y| + y^2 \ln |-y|] - 18[z^2 \ln |z| + z^2 \ln |-z|]|.
 \end{aligned}$$

Set  $x = -u, y = -v, z = -w$  it follows that

$$\begin{aligned}
 & |f_q(x + 2y + 3z) + f_q(x - 2y + 3z) + f_q(x + 2y - 3z) + f_q(x - 2y - 3z) \\
 & \quad - 4f_q(x) - 8[f_q(y) + f_q(-y)] - 18[f_q(z) + f_q(-z)]| \\
 & = |(-u - 2v - 3w)^2 \ln |-u - 2v - 3w| + (-u + 2v - 3w)^2 \ln |-u + 2v - 3w| \\
 & \quad + (-u - 2v + 3w)^2 \ln |-u - 2v + 3w| + (-u + 2v + 3w)^2 \ln |-u + 2v + 3w| \\
 & \quad + 4u^2 \ln |-u| - 8[v^2 \ln |-v| + v^2 \ln |v|] - 18[w^2 \ln |-w| + w^2 \ln |w|]|. \\
 & |f_q(-u - 2v - 3w) + f_q(-u + 2v - 3w) + f_q(-u - 2v + 3w) + f_q(-u + 2v + 3w) \\
 & \quad - 4f_q(-u) - 8[f_q(-v) + f_q(v)] - 18[f_q(-w) + f_q(w)]|. \\
 & \leq \lambda |-u|^{\frac{s}{3}} |-v|^{\frac{s}{3}} |-w|^{\frac{2-2s}{3}} = \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{2-2s}{3}}.
 \end{aligned}$$

**Case (iii):** If  $x > 0, y < 0, z < 0$  than  $x + 2z + 3z < 0, x - 2z + 3z < 0,$

$x + 2z - 3z < 0, x - 2z - 3z < 0$  and therefore (46) becomes,

$$\begin{aligned}
 & |f_q(x + 2y + 3z) + f_q(x - 2y + 3z) + f_q(x + 2y - 3z) + f_q(x - 2y - 3z) \\
 & \quad - 4f_q(x) - 8[f_q(y) + f_q(-y)] - 18[f_q(z) + f_q(-z)]|. \\
 & = |(x + 2y + 3z)^2 \ln |x + 2y + 3z| + (x - 2y + 3z)^2 \ln |x - 2y + 3z| + (x + 2y - 3z)^2 \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z)^2 \ln |x - 2y - 3z| - 4x^2 \ln |x| - 8[y^2 \ln |y| + y^2 \ln |-y|] - 18[z^2 \ln |z| + z^2 \ln |-z|]|.
 \end{aligned}$$

Set  $x = u, y = -v, z = -w$  it follows that

$$\begin{aligned}
 & |f_q(x + 2y + 3z) + f_q(x - 2y + 3z) + f_q(x + 2y - 3z) + f_q(x - 2y - 3z) \\
 & \quad - 4f_q(x) - 8[f_q(y) + f_q(-y)] - 18[f_q(z) + f_q(-z)]| \\
 & = |(u - 2v - 3w)^2 \ln |u - 2v - 3w| + (u + 2v - 3w)^2 \ln |u + 2v - 3w| + (u - 2v + 3w)^2 \ln |u - 2v + 3w| \\
 & \quad + (u + 2v + 3w)^2 \ln |u + 2v + 3w| - 4u^2 \ln |u| - 8[v^2 \ln |-v| + v^2 \ln |v|] - 18[w^2 \ln |-w| + w^2 \ln |w|]|. \\
 & |f_q(u - 2v - 3w) + f_q(u + 2v - 3w) + f_q(u - 2v + 3w) + f_q(u + 2v + 3w) \\
 & \quad - 4f_q(u) - 8[f_q(-v) + f_q(v)] - 18[f_q(-w) + f_q(w)]| \\
 & \leq \lambda |u|^{\frac{s}{3}} |-v|^{\frac{s}{3}} |-w|^{\frac{2-2s}{3}} = \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{2-2s}{3}}.
 \end{aligned}$$

**Case (iv):** If  $x < 0, y > 0, z > 0$  than  $x + 2z + 3z < 0, x - 2z + 3z < 0,$   
 $x + 2z - 3z < 0, x - 2z - 3z < 0$  and therefore (46) becomes,

$$\begin{aligned}
 & |f_q(x + 2y + 3z) + f_q(x - 2y + 3z) + f_q(x + 2y - 3z) + f_q(x - 2y - 3z) \\
 & \quad - 4f_q(x) - 8[f_q(y) + f_q(-y)] - 18[f_q(z) + f_q(-z)]| \\
 & = |(x + 2y + 3z)^2 \ln |x + 2y + 3z| + (x - 2y + 3z)^2 \ln |x - 2y + 3z| + (x + 2y - 3z)^2 \ln |x + 2y - 3z| \\
 & \quad + (x - 2y - 3z)^2 \ln |x - 2y - 3z| - 4x^2 \ln |x| - 8[y^2 \ln |y| + y^2 \ln |-y|] - 18[z^2 \ln |z| + z^2 \ln |-z|]|.
 \end{aligned}$$

Set  $x = -u, y = v, z = w$  it follows that

$$\begin{aligned}
 & |f_q(x + 2y + 3z) + f_q(x - 2y + 3z) + f_q(x + 2y - 3z) + f_q(x - 2y - 3z) \\
 & \quad - 4f_q(x) - 8[f_q(y) + f_q(-y)] - 18[f_q(z) + f_q(-z)]| \\
 & = |(-u + 2v + 3w)^2 \ln |-u + 2v + 3w| + (-u - 2v + 3w)^2 \ln |-u - 2v + 3w| \\
 & \quad + (-u + 2v - 3w)^2 \ln |-u + 2v - 3w| + (-u - 2v - 3w)^2 \ln |-u - 2v - 3w| \\
 & \quad + 4u^2 \ln |u| - 8[v^2 \ln |v| + v^2 \ln |-v|] - 18[w^2 \ln |w| + w^2 \ln |-w|]| \\
 & = |f_q(-u + 2v + 3w) + f_q(-u - 2v + 3w) + f_q(-u + 2v - 3w) + f_q(-u - 2v - 3w) \\
 & \quad - 4f_q(-u) - 8[f_q(v) + f_q(-v)] - 18[f_q(w) + f_q(-w)]| \\
 & \leq \lambda |-u|^{\frac{s}{3}} |v|^{\frac{s}{3}} |w|^{\frac{2-2s}{3}} = \lambda |x|^{\frac{s}{3}} |y|^{\frac{s}{3}} |z|^{\frac{2-2s}{3}}.
 \end{aligned}$$

**Case (v):** If  $x = y = z = 0$  in (46) then it is trivial. □

Now we provide an example to illustrate that the functional equation (6) is not stable for  $s = \frac{2}{3}$  in Condition (iii) of Corollary 3.7.

**Example 3.10.** Let  $\phi : \mathbb{R} \rightarrow \mathbb{R}$  be a function defined by

$$\phi(x) = \begin{cases} \mu x^2, & \text{if } |x| < \frac{2}{3}, \\ \frac{2\mu}{3}, & \text{otherwise} \end{cases}$$

where  $\mu > 0$  is a constant, and define a function  $f_q : \mathbb{R} \rightarrow \mathbb{R}$  by

$$f_q(x) = \sum_{n=0}^{\infty} \frac{\phi(6^n x)}{36^n} \quad \text{for all } x \in \mathbb{R}.$$

Then  $f_q$  satisfies the functional inequality

$$|Df_q(x, y, z)| \leq \frac{1440}{35} \mu \left( |x|^{\frac{2}{3}} |y|^{\frac{2}{3}} |z|^{\frac{2}{3}} + |x|^2 + |y|^2 + |z|^2 \right), \quad (69)$$

for all  $x, y, z \in \mathbb{R}$ . Then there doesn't exist a quadratic mapping  $Q : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\beta > 0$  such that

$$|f_q(x) - Q(x)| \leq \beta |x|^2 \quad \text{for all } x \in \mathbb{R}. \quad (70)$$

*Proof.* Now

$$|f_q(x)| \leq \sum_{n=0}^{\infty} \frac{|\phi(6^n x)|}{|36^n|} = \sum_{n=0}^{\infty} \frac{\mu}{3} \times \frac{1}{36^n} = \frac{2\mu}{5}.$$

Therefore we see that  $f_q$  is bounded. We are going to prove that  $f_q$  satisfies (69).

If  $x = y = z = 0$ , then (69) is trivial. If  $|x|^{\frac{2}{3}} |y|^{\frac{2}{3}} |z|^{\frac{2}{3}} + |x|^2 + |y|^2 + |z|^2 \geq \frac{1}{36}$  then the left hand side of (48) is less than  $\frac{1440}{35} \mu$ . Now suppose that  $0 < |x|^{\frac{2}{3}} |y|^{\frac{2}{3}} |z|^{\frac{2}{3}} + |x|^2 + |y|^2 + |z|^2 < \frac{1}{36}$ . Then there exists a positive integer  $k$  such that

$$\frac{1}{36^{k+2}} \leq |x|^{\frac{2}{3}} |y|^{\frac{2}{3}} |z|^{\frac{2}{3}} + |x|^2 + |y|^2 + |z|^2 < \frac{1}{36^{k+1}}, \quad (71)$$

so that  $6^{k-1} |x|^{\frac{2}{3}} |y|^{\frac{2}{3}} |z|^{\frac{2}{3}} < \frac{1}{6}$ ,  $6^{k-1} |x| < \frac{1}{6}$ ,  $6^{k-1} |y| < \frac{1}{6}$ ,  $6^{k-1} |z| < \frac{1}{6}$  and consequently

$$\begin{aligned} &6^{k-1}(x + 2y + 3z), 6^{k-1}(x - 2y + 3z), 6^{k-1}(x + 2y - 3z), 6^{k-1}(x - 2y - 3z), \\ &6^{k-1}(x), 6^{k-1}(-x), 6^{k-1}(y), 6^{k-1}(-y), 6^{k-1}(z), 6^{k-1}(-z) \in \left( \frac{-1}{6}, \frac{1}{6} \right). \end{aligned}$$

Therefore for each  $n = 0, 1, \dots, k - 1$ , we have

$$\begin{aligned} &6^n(x + 2y + 3z), 6^n(x - 2y + 3z), 6^n(x + 2y - 3z), 6^n(x - 2y - 3z), \\ &6^n(x), 6^n(-x), 6^n(y), 6^n(-y), 6^n(z), 6^n(-z) \in \left( \frac{-1}{6}, \frac{1}{6} \right) \end{aligned}$$

and

$$\begin{aligned} &\phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \\ &- 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] = 0 \end{aligned}$$

for  $n = 0, 1, \dots, k - 1$ . From the definition of  $f_q$  and (71), we obtain that

$$\begin{aligned} &\left| Df_q(x, y, z) \right| \\ &= \sum_{n=0}^{\infty} \frac{1}{36^n} \left| \phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \right. \\ &\quad \left. - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] \right| \\ &= \sum_{n=k}^{\infty} \frac{1}{36^n} \left| \phi(6^n(x + 2y + 3z)) + \phi(6^n(x - 2y + 3z)) + \phi(6^n(x + 2y - 3z)) + \phi(6^n(x - 2y - 3z)) \right. \\ &\quad \left. - 4\phi(6^n x) - 8[\phi(6^n y) + \phi(-6^n y)] - 18[\phi(6^n z) + \phi(-6^n z)] \right| \\ &\leq \sum_{n=k}^{\infty} \frac{1}{36^n} \frac{120}{3} \mu = \frac{120}{3} \mu \times \frac{1}{36^k} \times \frac{36}{35} \leq \frac{1440}{35} \mu \left( |x|^{\frac{2}{3}} |y|^{\frac{2}{3}} |z|^{\frac{2}{3}} + |x|^2 + |y|^2 + |z|^2 \right). \end{aligned}$$

Thus  $f_q$  satisfies (69) for all  $x, y, z \in \mathbb{R}$  with  $0 < |x|^{\frac{2}{3}}|y|^{\frac{2}{3}}|z|^{\frac{2}{3}} + |x|^2 + |y|^2 + |z|^2 < \frac{1}{6}$ .

We claim that the additive functional equation (6) is not stable for  $s = \frac{2}{3}$  in condition (iv) of Corollary 3.2. Suppose on the contrary that there exists an additive mapping  $Q : \mathbb{R} \rightarrow \mathbb{R}$  and a constant  $\beta > 0$  satisfying (70). Since  $f_q$  is bounded and continuous for all  $x \in \mathbb{R}$ ,  $Q$  is bounded on any open interval containing the origin and continuous at the origin. In view of Theorem 3.1,  $A$  must have the form  $Q(x) = cx^2$  for any  $x$  in  $\mathbb{R}$ . Thus we obtain that

$$|f_q(x)| \leq (\beta + |c|)|x|^2. \quad (72)$$

But we can choose a positive integer  $m$  with  $m\mu > \beta + |c|$ .

If  $x \in (0, \frac{1}{6^{m-1}})$ , then  $6^n x \in (0, \frac{1}{6})$  for all  $n = 0, 1, \dots, m-1$ . For this  $x$ , we get

$$f_q(x) = \sum_{n=0}^{\infty} \frac{\alpha(6^n x)}{36^n} \geq \sum_{n=0}^{m-1} \frac{\mu(6^n x)^2}{36^n} = m\mu x^2 > (\beta + |c|)x^2,$$

which contradicts (72). Therefore the quadratic functional equation (6) is not stable in the sense of Ulam, Hyers and Rassias if  $s = \frac{2}{3}$ , assumed in the inequality (40).  $\square$

**Theorem 3.11.** Let  $j \in \{-1, 1\}$  and  $\alpha, \beta : X^3 \rightarrow [0, \infty)$  be a function satisfying (28) and (52) for all  $x, y, z \in X$ . Let  $f : X \rightarrow Y$  be a function satisfying the inequality

$$\|Df(x, y, z)\| \leq \alpha(x, y, z) \quad (73)$$

for all  $x, y, z \in X$ . Then there exists a unique additive mapping  $A : X \rightarrow Y$  and a unique quadratic mapping  $Q : X \rightarrow Y$  satisfying (6) and

$$\|f(x) - A(x) - Q(x)\| \leq \frac{1}{2} \left[ \frac{1}{12} \sum_{k=\frac{1-j}{2}}^{\infty} \left( \frac{\beta(6^{kj}x) + \beta(-6^{kj}x)}{6^{kj}} \right) + \frac{1}{72} \sum_{k=\frac{1-j}{2}}^{\infty} \left( \frac{\beta(6^{kj}x) + \beta(-6^{kj}x)}{36^{kj}} \right) \right], \quad (74)$$

where  $\beta(6^{kj}x)$ ,  $A(x)$  and  $Q(x)$  are defined in (31), (55), (32) and (56), respectively, for all  $x \in X$ .

*Proof.* Let  $f_o(x) = \frac{f_a(x) - f_a(-x)}{2}$  for all  $x \in X$ . Then  $f_o(0) = 0$  and  $f_o(-x) = -f_o(x)$  for all  $x \in X$ . Hence

$$\|Df_o(x, y, z)\| \leq \frac{\alpha(x, y, z)}{2} + \frac{\alpha(-x, -y, -z)}{2} \quad (75)$$

for all  $x, y, z \in X$ . By Theorem 3.1, we have

$$\|f_o(x) - A(x)\| \leq \frac{1}{24} \sum_{k=\frac{1-j}{2}}^{\infty} \left( \frac{\beta(6^{kj}x) + \beta(-6^{kj}x)}{6^{kj}} \right) \quad (76)$$

for all  $x \in X$ . Also, let  $f_e(x) = \frac{f_q(x) + f_q(-x)}{2}$  for all  $x \in X$ . Then  $f_e(0) = 0$  and  $f_e(-x) = f_e(x)$  for all  $x \in X$ . Hence

$$\|Df_e(x, y, z)\| \leq \frac{\alpha(x, y, z)}{2} + \frac{\alpha(-x, -y, -z)}{2} \quad (77)$$

for all  $x, y, z \in X$ . By Theorem 3.6, we have

$$\|f_e(x) - Q(x)\| \leq \frac{1}{144} \sum_{k=\frac{1-j}{2}}^{\infty} \left( \frac{\beta(6^{kj}x) + \beta(-6^{kj}x)}{36^{kj}} \right) \tag{78}$$

for all  $x \in X$ . Define

$$f(x) = f_e(x) + f_o(x) \tag{79}$$

for all  $x \in X$ . From (76),(78) and (79), we arrive

$$\begin{aligned} \|f(x) - A(x) - Q(x)\| &= \|f_e(x) + f_o(x) - A(x) - Q(x)\| \\ &\leq \|f_o(x) - A(x)\| + \|f_e(x) - Q(x)\| \\ &\leq \frac{1}{2} \left[ \frac{1}{12} \sum_{k=\frac{1-j}{2}}^{\infty} \left( \frac{\beta(6^{kj}x) + \beta(-6^{kj}x)}{6^{kj}} \right) \right. \\ &\quad \left. + \frac{1}{72} \sum_{k=\frac{1-j}{2}}^{\infty} \left( \frac{\beta(6^{kj}x) + \beta(-6^{kj}x)}{36^{kj}} \right) \right] \end{aligned}$$

for all  $x \in X$ . Hence the theorem is proved. □

Using Corollaries 3.2 and 3.7, we have the following corollary concerning the stability of (6).

**Corollary 3.12.** *Let  $\lambda$  and  $s$  be nonnegative real numbers. Let a function  $f : X \rightarrow Y$  satisfy the inequality*

$$\|Df(x, y, z)\| \leq \begin{cases} \lambda, & s \neq 1, 2; \\ \lambda \{ \|x\|^s + \|y\|^s + \|z\|^s \}, & s \neq 1, 2; \\ \lambda \|x\|^s \|y\|^s \|z\|^s, & 3s \neq 1, 2; \\ \lambda \{ \|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \} \}, & 3s \neq 1, 2; \end{cases} \tag{80}$$

for all  $x, y, z \in X$ . Then there exists a unique additive function  $A : X \rightarrow Y$  and a unique quadratic function  $Q : X \rightarrow Y$  such that

$$\|f(x) - A(x) - Q(x)\| \leq \begin{cases} 3\lambda \left( \frac{1}{10} + \frac{1}{70} \right), \\ 4\lambda \|x\|^s \left( \frac{1}{|6 - 6^s|} + \frac{1}{|36 - 6^s|} \right), \\ \lambda \|x\|^{3s} \left( \frac{1}{|6 - 6^{3s}|} + \frac{1}{|36 - 6^{3s}|} \right), \\ 5\lambda \|x\|^{3s} \left( \frac{1}{|6 - 6^{3s}|} + \frac{1}{|36 - 6^{3s}|} \right), \end{cases} \tag{81}$$

for all  $x \in X$ .

### 4. Stability Results: Fixed Point Method

In this section, the authors have proved the generalized Ulam - Hyers stability of functional equation (6) in Banach spaces with the help of the fixed point method. Now we will recall the fundamental result in the fixed point theory.

**Theorem 4.1.** [24] *(The alternative of fixed point)* Suppose that for a complete generalized metric space  $(X, d)$  and a strictly contractive mapping  $T : X \rightarrow X$  with Lipschitz constant  $L$ . Then, for each given element  $x \in X$ , either

$$(B_1) \quad d(T^n x, T^{n+1} x) = \infty \quad \forall \quad n \geq 0,$$

or

(B<sub>2</sub>) there exists a natural number  $n_0$  such that:

$$(i) \quad d(T^n x, T^{n+1} x) < \infty \text{ for all } n \geq n_0 ;$$

(ii) The sequence  $(T^n x)$  is convergent to a fixed point  $y^*$  of  $T$ ;

(iii)  $y^*$  is the unique fixed point of  $T$  in the set  $Y = \{y \in X : d(T^{n_0} x, y) < \infty\}$ ;

$$(iv) \quad d(y^*, y) \leq \frac{1}{1-L} d(y, Ty) \text{ for all } y \in Y.$$

Hereafter throughout this section, let us assume  $V$  be a vector space and  $B$  Banach space respectively. Define a mapping  $Df : V \rightarrow B$  by

$$\begin{aligned} Df(x, y, z) &= f(x + 2y + 3z) + f(x - 2y + 3z) + f(x + 2y - 3z) + f(x - 2y - 3z) \\ &\quad - 4f(x) - 8[f(y) + f(-y)] - 18[f(z) + f(-z)] \end{aligned}$$

for all  $x, y, z \in V$ .

**Theorem 4.2.** Let  $f_a : V \rightarrow B$  be a mapping for which there exists functions  $\alpha, \beta, \gamma : V^3 \rightarrow [0, \infty)$  with the condition

$$\lim_{k \rightarrow \infty} \frac{\alpha(\mu_i^k x, \mu_i^k y, \mu_i^k z)}{\mu_i^k} = 0, \quad (82)$$

where

$$\mu_i = \begin{cases} 6, & i = 0, \\ \frac{1}{6}, & i = 1 \end{cases}$$

satisfying the functional inequality

$$\|Df_a(x, y, z)\| \leq \alpha(x, y, z) \quad (83)$$

for all  $x, y, z \in V$ . If there exists an  $L = L(i) < 1$  such that the function

$$x \rightarrow \gamma(x) = \frac{1}{2} \beta \left( \frac{x}{6} \right),$$

one has the property

$$\gamma(x) = L \mu_i \gamma \left( \frac{x}{\mu_i} \right) \quad (84)$$

for all  $x \in V$ . Then there exists a unique additive function  $A : V \rightarrow B$  satisfying the functional equation (6) and

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) \quad (85)$$

holds for all  $x \in V$ .

*Proof.* Consider the set  $X = \{p/p : V \rightarrow B, p(0) = 0\}$  and introduce the generalized metric on  $X$ ,

$$d(p, q) = \inf\{K \in (0, \infty) : \|p(x) - q(x)\| \leq K\gamma(x), x \in V\}.$$

It is easy to see that  $(X, d)$  is complete.

Define  $T : X \rightarrow X$  by

$$Tp(x) = \frac{1}{\mu_i}p(\mu_i x), \forall x \in V.$$

Now  $p, q \in X$ ,

$$\begin{aligned} d(p, q) \leq K &\Rightarrow \|p(x) - q(x)\| \leq K\gamma(x), x \in V. \\ &\Rightarrow \left\| \frac{1}{\mu_i}p(\mu_i x) - \frac{1}{\mu_i}q(\mu_i x) \right\| \leq \frac{1}{\mu_i}K\gamma(\mu_i x), x \in V, \\ &\Rightarrow \left\| \frac{1}{\mu_i}p(\mu_i x) - \frac{1}{\mu_i}q(\mu_i x) \right\| \leq LK\gamma(x), x \in V, \\ &\Rightarrow \|Tp(x) - Tq(x)\| \leq LK\gamma(x), x \in V, \\ &\Rightarrow d(Tp, Tq) \leq LK. \end{aligned}$$

This implies

$$d(Tp, Tq) \leq Ld(p, q),$$

for all  $p, q \in X$ . i.e.,  $T$  is a strictly contractive mapping on  $X$  with Lipschitz constant  $L$ .

From (36), we have

$$\left\| \frac{f_a(6x)}{6} - f_a(x) \right\| \leq \frac{\beta(x)}{12} \tag{86}$$

where

$$\beta(x) = 2\alpha(x, x, x) + \alpha(x, 0, x)$$

for all  $x \in V$ . Using (84) for the case  $i = 0$ , it reduces to

$$\left\| \frac{1}{6}f_a(6x) - f_a(x) \right\| \leq \frac{1}{6}\gamma(x)$$

for all  $x \in V$ .

$$\text{i.e., } d(Tf_a, f_a) \leq \frac{1}{6} = L = L^{1-0} = L^{1-i} < \infty.$$

Again replacing  $x = \frac{x}{6}$  in (86), we get

$$\left\| f_a(x) - 6f\left(\frac{x}{6}\right) \right\| \leq \frac{1}{2}\beta\left(\frac{x}{6}\right).$$

for all  $x \in V$ . Using (84) for the case  $i = 1$ , it reduces to

$$\left\| f_a(x) - 6f\left(\frac{x}{6}\right) \right\| \leq \gamma(x)$$

for all  $x \in V$ .

$$\text{i.e., } d(f_a, Tf_a) \leq 1 = L^0 = L^{1-1} = L^{1-i} < \infty.$$

In the above cases, we arrive

$$d(f_a, Tf_a) \leq L^{1-i}.$$

Therefore  $(B_2(i))$  holds.

By  $(B_2(ii))$ , it follows that there exists a fixed point  $A$  of  $T$  in  $X$  such that

$$A(x) = \lim_{k \rightarrow \infty} \frac{f_a(\mu_i^k x)}{\mu_i^k}, \quad \forall x \in V. \quad (87)$$

Claim that  $A : V \rightarrow B$  is additive. Replacing  $(x, y, z)$  by  $(\mu_i^k x, \mu_i^k y, \mu_i^k z)$  in (83) and dividing by  $\mu_i^k$ , it follows from (82) and (87),  $A$  satisfies (6) for all  $x, y, z \in V$ .

By  $(B_2(iii))$ ,  $A$  is the unique fixed point of  $T$  in the set  $Y = \{f_a \in X : d(Tf_a, A) < \infty\}$ , using the fixed point alternative result  $A$  is the unique function such that

$$\|f_a(x) - A(x)\| \leq K\gamma(x)$$

for all  $x \in V$  and  $K > 0$ . Finally by  $(B_2(iv))$ , we obtain

$$d(f_a, A) \leq \frac{1}{1-L} d(f_a, Tf_a)$$

implying

$$d(f_a, A) \leq \frac{L^{1-i}}{1-L}.$$

Hence we conclude that

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x).$$

for all  $x \in V$ . This completes the proof of the theorem.  $\square$

From Theorem 4.2, we obtain the following corollary concerning the stability for the functional equation (6).

**Corollary 4.3.** *Let  $f_a : V \rightarrow B$  be a mapping and there exist real numbers  $\lambda$  and  $s$  such that*

$$\|Df_a(x, y, z)\| \leq \begin{cases} (i) & \lambda, \\ (ii) & \lambda \{|x|^s + |y|^s + |z|^s\}, & s < 1 \text{ or } s > 1; \\ (iii) & \lambda |x|^s |y|^s |z|^s, & 3s < 1 \text{ or } 3s > 1; \\ (iv) & \lambda \{|x|^s |y|^s |z|^s + \{|x|^{3s} + |y|^{3s} + |z|^{3s}\}\}, & 3s < 1 \text{ or } 3s > 1; \end{cases} \quad (88)$$

for all  $x, y, z \in V$ , then there exists a unique additive function  $A : V \rightarrow B$  such that

$$\|f_a(x) - A(x)\| \leq \begin{cases} (i) & \frac{3\lambda}{10}, \\ (ii) & \frac{4\lambda|x|^s}{|6-6^s|}, \\ (iii) & \frac{\lambda|x|^{3s}}{|6-6^{3s}|}, \\ (iv) & \frac{5\lambda|x|^{3s}}{|6-6^{3s}|} \end{cases} \quad (89)$$

for all  $x \in V$ .



*Proof.* Let us set

$$\alpha(x, y, z) = \begin{cases} \lambda, \\ \lambda \{ \|x\|^s + \|y\|^s + \|z\|^s \}, \\ \lambda \|x\|^s \|y\|^s \|z\|^s, \\ \lambda \{ \|x\|^s \|y\|^s \|z\|^s + (\|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s}) \} \end{cases}$$

for all  $x, y, z \in V$ . Now

$$\begin{aligned} \frac{\alpha(\mu_i^k x, \mu_i^k y, \mu_i^k z)}{\mu_i^k} &= \begin{cases} \frac{\lambda}{\mu_i^k}, \\ \frac{\lambda}{\mu_i^k} \{ \|\mu_i^k x\|^s + \|\mu_i^k y\|^s + \|\mu_i^k z\|^s \}, \\ \frac{\lambda}{\mu_i^k} \|\mu_i^k x\|^s \|\mu_i^k y\|^s \|\mu_i^k z\|^s, \\ \frac{\lambda}{\mu_i^k} \{ \|\mu_i^k x\|^s \|\mu_i^k y\|^s \|\mu_i^k z\|^s + (\|\mu_i^k x\|^{3s} + \|\mu_i^k y\|^{3s} + \|\mu_i^k z\|^{3s}) \} \end{cases} \\ &= \begin{cases} \rightarrow 0 \text{ as } k \rightarrow \infty, \\ \rightarrow 0 \text{ as } k \rightarrow \infty, \\ \rightarrow 0 \text{ as } k \rightarrow \infty, \\ \rightarrow 0 \text{ as } k \rightarrow \infty. \end{cases} \end{aligned}$$

i.e., (82) is holds. But, we have

$$\gamma(x) = \frac{1}{2} \beta\left(\frac{x}{6}\right) = \frac{1}{2} \left[ 2\alpha\left(\frac{x}{6}, \frac{x}{6}, \frac{x}{6}\right) + \alpha\left(\frac{x}{6}, 0, \frac{x}{6}\right) \right].$$

Hence

$$\gamma(x) = \frac{1}{2} \left[ 2\alpha\left(\frac{x}{6}, \frac{x}{6}, \frac{x}{6}\right) + \alpha\left(\frac{x}{6}, 0, \frac{x}{6}\right) \right] = \begin{cases} \frac{3\lambda}{2}, \\ \frac{4\lambda}{6^s} \|x\|^s, \\ \frac{\lambda}{6^{3s}} \|x\|^{3s}, \\ \frac{5\lambda}{6^{3s}} \|x\|^{3s}. \end{cases}$$

Also,

$$\frac{1}{\mu_i} \gamma(\mu_i x) = \begin{cases} \frac{3\lambda}{\mu_i \cdot 2}, \\ \frac{4\lambda}{\mu_i \cdot 6^s} \|\mu_i x\|^s, \\ \frac{\lambda}{\mu_i \cdot 6^{3s}} \|\mu_i x\|^{3s}, \\ \frac{5\lambda}{\mu_i \cdot 6^{3s}} \|\mu_i x\|^{3s}. \end{cases} = \begin{cases} \mu_i^{-1} \frac{3\lambda}{2}, \\ \mu_i^{s-1} \frac{4\lambda}{6^s} \|x\|^s, \\ \mu_i^{3s-1} \frac{\lambda}{6^{3s}} \|x\|^{3s}, \\ \mu_i^{3s-1} \frac{5\lambda}{6^{3s}} \|x\|^{3s}. \end{cases} = \begin{cases} \mu_i^{-1} \gamma(x), \\ \mu_i^{s-1} \gamma(x), \\ \mu_i^{3s-1} \gamma(x), \\ \mu_i^{3s-1} \gamma(x). \end{cases}$$

Hence the inequality (84) holds either,  $L = 6^{-1}$  for  $s = 1$  if  $i = 0$ , or  $L = 6$  for  $s = 0$  if  $i = 1$ . Now from (85), we prove the following cases for condition (i).

**Case:1**  $L = 6^{-1}$  for  $s = 1$  if  $i = 0$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) = \frac{(6^{-1})^{1-0}}{1-(6^{-1})} \cdot \frac{3\lambda}{2} = \frac{3\lambda}{10}.$$

**Case:2**  $L = 6$  for  $s = 0$  if  $i = 1$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) = \frac{(6)^{1-1}}{1-6} \cdot \frac{3\lambda}{2} = \frac{-3\lambda}{10}.$$

Again, (84) holds either,  $L = 6^{s-1}$  for  $s < 1$  if  $i = 0$ , or  $L = \frac{1}{6^{s-1}}$  for  $s > 1$  if  $i = 1$ . Now from (85), we prove the following cases for condition (ii).

**Case:1**  $L = 6^{s-1}$  for  $s < 1$  if  $i = 0$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) = \frac{(6^{s-1})^{1-0}}{1-6^{s-1}} \frac{4\lambda}{6^s} \|x\|^s = \frac{6^s}{6-6^s} \frac{4\lambda}{6^s} \|x\|^s = \frac{4\lambda \|x\|^s}{6-6^s}.$$

**Case:2**  $L = \frac{1}{6^{s-1}}$  for  $s > 1$  if  $i = 1$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) = \frac{\left(\frac{1}{6^{s-1}}\right)^{1-1}}{1-\frac{1}{6^{s-1}}} \frac{4\lambda}{6^s} \|x\|^s = \frac{6^s}{6^s-6} \frac{4\lambda}{6^s} \|x\|^s = \frac{4\lambda \|x\|^s}{6^s-6}.$$

Also, (84) holds either,  $L = 6^{3s-1}$  for  $3s < 1$  if  $i = 0$ , or  $L = \frac{1}{6^{3s-1}}$  for  $3s > 1$  if  $i = 1$ . Now from (85), we prove the following cases for condition (iii).

**Case:1**  $L = 6^{3s-1}$  for  $3s < 1$  if  $i = 0$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) = \frac{(6^{3s-1})^{1-0}}{1-6^{3s-1}} \frac{\lambda}{6^{3s}} \|x\|^{3s} = \frac{6^{3s}}{6-6^{3s}} \frac{\lambda}{6^{3s}} \|x\|^{3s} = \frac{\lambda \|x\|^{3s}}{6-6^{3s}}.$$

**Case:2**  $L = \frac{1}{6^{3s-1}}$  for  $3s > 1$  if  $i = 1$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) = \frac{\left(\frac{1}{6^{3s-1}}\right)^{1-1}}{1-\frac{1}{6^{3s-1}}} \frac{\lambda}{6^{3s}} \|x\|^{3s} = \frac{6^{3s}}{6^{3s}-6} \frac{\lambda}{6^{3s}} \|x\|^{3s} = \frac{\lambda \|x\|^{3s}}{6^{3s}-6}.$$

Finally, (84) holds either,  $L = 6^{3s-1}$  for  $3s < 1$  if  $i = 0$ , or  $L = \frac{1}{6^{3s-1}}$  for  $3s > 1$  if  $i = 1$ . Now from (85), we prove the following cases for condition (iv).

**Case:1**  $L = 6^{3s-1}$  for  $3s < 1$  if  $i = 0$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) = \frac{(6^{3s-1})^{1-0}}{1-6^{3s-1}} \frac{5\lambda}{6^{3s}} \|x\|^{3s} = \frac{6^{3s}}{6-6^{3s}} \frac{5\lambda}{6^{3s}} \|x\|^{3s} = \frac{5\lambda \|x\|^{3s}}{6-6^{3s}}.$$

**Case:2**  $L = \frac{1}{6^{3s-1}}$  for  $3s > 1$  if  $i = 1$

$$\|f_a(x) - A(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) = \frac{\left(\frac{1}{6^{3s-1}}\right)^{1-1}}{1-\frac{1}{6^{3s-1}}} \frac{5\lambda}{6^{3s}} \|x\|^{3s} = \frac{6^{3s}}{6^{3s}-6} \frac{5\lambda}{6^{3s}} \|x\|^{3s} = \frac{5\lambda \|x\|^{3s}}{6^{3s}-6}.$$

Hence the proof of the corollary. □

The proofs of the following Theorem and Corollary are similar to those proofs of Theorem 4.2 and Corollary 4.3 using (60). Hence we omit the proofs.

**Theorem 4.4.** Let  $f_q : V \rightarrow B$  be a mapping for which there exists functions  $\alpha, \beta, \gamma : V^3 \rightarrow [0, \infty)$  with the condition

$$\lim_{k \rightarrow \infty} \frac{\alpha(\mu_i^k x, \mu_i^k y, \mu_i^k z)}{\mu_i^{2k}} = 0 \tag{90}$$

where

$$\mu_i = \begin{cases} 6, & i = 0, \\ \frac{1}{6}, & i = 1 \end{cases}$$

satisfying the functional inequality

$$\|Df_q(x, y, z)\| \leq \alpha(x, y, z) \tag{91}$$

for all  $x, y, z \in V$ . If there exists  $L = L(i) < 1$  such that the function

$$x \rightarrow \gamma(x) = \frac{1}{2}\beta\left(\frac{x}{6}\right),$$

has the property

$$\gamma(x) = L \mu_i^2 \gamma\left(\frac{x}{\mu_i}\right) \tag{92}$$

for all  $x \in V$ . Then there exists a unique quadratic function  $Q : V \rightarrow B$  satisfying the functional equation (6) and

$$\|f_q(x) - Q(x)\| \leq \frac{L^{1-i}}{1-L} \gamma(x) \tag{93}$$

holds for all  $x \in V$ .

**Corollary 4.5.** Let  $f_q : V \rightarrow B$  be a mapping and there exist real numbers  $\lambda$  and  $s$  such that

$$\|Df_q(x, y, z)\| \leq \begin{cases} (i) & \lambda, \\ (ii) & \lambda \{ \|x\|^s + \|y\|^s + \|z\|^s \}, & s < 2 \text{ or } s > 2; \\ (iii) & \lambda \|x\|^s \|y\|^s \|z\|^s, & 3s < 2 \text{ or } 3s > 2; \\ (iv) & \lambda \{ \|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \} \}, & 3s < 2 \text{ or } 3s > 2; \end{cases} \tag{94}$$

for all  $x, y, z \in V$ , then there exists a unique quadratic function  $Q : V \rightarrow B$  such that

$$\|f_q(x) - Q(x)\| \leq \begin{cases} (i) & \frac{3\lambda}{70}, \\ (ii) & \frac{4\lambda \|x\|^s}{|36 - 6^s|}, \\ (iii) & \frac{\lambda \|x\|^{3s}}{|36 - 6^{3s}|} \\ (iv) & \frac{5\lambda \|x\|^{3s}}{|36 - 6^{3s}|} \end{cases} \tag{95}$$

for all  $x \in V$ .

**Theorem 4.6.** Let  $f : V \rightarrow B$  be a mapping for which there exists functions  $\alpha, \beta, \gamma : V^3 \rightarrow [0, \infty)$  with the condition (82) and (90), where

$$\mu_i = \begin{cases} 6, & i = 0, \\ \frac{1}{6}, & i = 1 \end{cases}$$

such that the functional inequality

$$\|Df(x, y, z)\| \leq \alpha(x, y, z) \tag{96}$$

holds for all  $x, y, z \in V$ . Assume there exists  $L = L(i) < 1$  such that the function

$$x \rightarrow \gamma(x) = \frac{1}{2}\beta\left(\frac{x}{6}\right),$$

has the properties (84) and (92) for all  $x \in V$ . Then there exists a unique additive function  $A : V \rightarrow B$  and a unique quadratic function  $Q : V \rightarrow B$  satisfying the functional equation (6) and

$$\|f(x) - A(x) - Q(x)\| \leq \frac{L^{1-i}}{1-L} (\gamma(x) + \gamma(-x)) \quad (97)$$

holds for all  $x \in V$ .

*Proof.* Let  $f_o(x) = \frac{f_a(x) - f_a(-x)}{2}$  for all  $x \in V$ . Then  $f_o(0) = 0$  and  $f_o(-x) = -f_o(x)$  for all  $x \in V$ . Hence

$$\|Df_o(x, y, z)\| \leq \frac{\alpha(x, y, z)}{2} + \frac{\alpha(-x, -y, -z)}{2}, \quad (98)$$

for all  $x, y, z \in V$ . By Theorem 4.2, we have

$$\|f_o(x) - A(x)\| \leq \frac{1}{2} \frac{L^{1-i}}{1-L} (\gamma(x) + \gamma(-x)), \quad (99)$$

for all  $x \in V$ . Also, let  $f_e(x) = \frac{f_q(x) + f_q(-x)}{2}$  for all  $x \in V$ . Then  $f_e(0) = 0$  and  $f_e(-x) = f_e(x)$  for all  $x \in V$ . Hence

$$\|Df_e(x, y, z)\| \leq \frac{\alpha(x, y, z)}{2} + \frac{\alpha(-x, -y, -z)}{2}, \quad (100)$$

for all  $x, y, z \in V$ . By Theorem 4.4, we have

$$\|f_e(x) - Q(x)\| \leq \frac{1}{2} \frac{L^{1-i}}{1-L} (\gamma(x) + \gamma(-x)), \quad (101)$$

for all  $x \in V$ . Define

$$f(x) = f_e(x) + f_o(x) \quad (102)$$

for all  $x \in V$ . From (99), (101) and (102), we arrive

$$\begin{aligned} \|f(x) - A(x) - Q(x)\| &= \|f_e(x) + f_o(x) - A(x) - Q(x)\| \\ &\leq \|f_o(x) - A(x)\| + \|f_e(x) - Q(x)\| \\ &\leq \frac{1}{2} \frac{L^{1-i}}{1-L} (\gamma(x) + \gamma(-x)) + \frac{1}{2} \frac{L^{1-i}}{1-L} (\gamma(x) + \gamma(-x)) \\ &\leq \frac{L^{1-i}}{1-L} (\gamma(x) + \gamma(-x)) \end{aligned}$$

for all  $x \in V$ . Hence the theorem is proved.  $\square$

Using Corollaries 4.3 and 4.5, we have the following corollary concerning the stability of (6).

**Corollary 4.7.** Let  $\lambda$  and  $s$  be nonnegative real numbers. Let a function  $f : V \rightarrow B$  satisfy the inequality

$$\|Df(x, y, z)\| \leq \begin{cases} (i) & \lambda, \\ (ii) & \lambda \{ \|x\|^s + \|y\|^s + \|z\|^s \}, & s \neq 1, 2; \\ (iii) & \lambda \|x\|^s \|y\|^s \|z\|^s, & 3s \neq 1, 2; \\ (iv) & \lambda \{ \|x\|^s \|y\|^s \|z\|^s + \{ \|x\|^{3s} + \|y\|^{3s} + \|z\|^{3s} \} \}, & 3s \neq 1, 2; \end{cases} \quad (103)$$

for all  $x, y, z \in V$ . Then there exists a unique additive function  $A : V \rightarrow B$  and a unique quadratic function  $Q : V \rightarrow B$  such that

$$\|f(x) - A(x) - Q(x)\| \leq \begin{cases} (i) & 3\lambda \left( \frac{1}{10} + \frac{1}{70} \right), \\ (ii) & 4\lambda \|x\|^s \left( \frac{1}{|6 - 6^s|} + \frac{1}{|36 - 6^s|} \right), \\ (iii) & \lambda \|x\|^{3s} \left( \frac{1}{|6 - 6^{3s}|} + \frac{1}{|36 - 6^{3s}|} \right), \\ (iv) & 5\lambda \|x\|^{3s} \left( \frac{1}{|6 - 6^{3s}|} + \frac{1}{|36 - 6^{3s}|} \right), \end{cases} \quad (104)$$

for all  $x \in V$ .

## References

- [1] J.Aczel and J.Dhombres, *Functional Equations in Several Variables*, Cambridge Univ, Press, (1989).
- [2] T.Aoki, *On the stability of the linear transformation in Banach spaces*, J. Math. Soc. Japan, 2(1950), 64-66.
- [3] M.Arunkumar and S. Karthikeyan, *Solution and stability of n-dimensional mixed Type additive and quadratic functional equation*, Far East Journal of Applied Mathematics, 54(1)(2011), 47-64.
- [4] M.Arunkumar, S.Hema latha and C.Devi Shaymala Mary, *Functional equation originating from arithmetic Mean of consecutive terms of an arithmetic Progression are stable in banach space: Direct and fixed point method*, JP Journal of Mathematical Sciences, 3(1)(2012), 27-43.
- [5] M.Arunkumar, Matina J.Rassias and Yanhui Zhang, *Ulam - Hyers stability of a 2- variable AC - mixed type functional equation: direct and fixed point methods*, Journal of Modern Mathematics Frontier (JMMF), 1(3)(2012), 10-26.
- [6] M.Arunkumar and John M.Rassias, *On the generalized Ulam-Hyers stability of an AQ-mixed type functional equation with counter examples*, Far East Journal of Applied Mathematics, 71(2)(2012), 279-305.
- [7] M.Arunkumar and P.Agilan, *Additive Quadratic functional equation are Stable in Banach space: A Fixed Point Approach*, International Journal of pure and Applied Mathematics, 86(6)(2013), 951-963.
- [8] M.Arunkumar and P.Agilan, *Additive Quadratic functional equation are Stable in Banach space: A Direct Method*, Far East Journal of Mathematical Sciences, 80(1)(2013), 105-121.
- [9] I.S.Chang, E.H.Lee and H.M.Kim, *On the Hyers-Ulam-Rassias stability of a quadratic functional equations*, Math. Ineq. Appl., 6(1)(2003), 87-95.
- [10] S.Czerwik, *Functional Equations and Inequalities in Several Variables*, World Scientific, River Edge, NJ, (2002).
- [11] P.Gavruta, *A generalization of the Hyers-Ulam-Rassias stability of approximately additive mappings*, J. Math. Anal. Appl., 184(1994), 431-436.
- [12] M.Eshaghi Gordji, N.Ghobadipour and J. M.Rassias, *Fuzzy stability of additive-quadratic functional Equations*, arXiv:0903.0842v1 [math.FA] 4 Mar 2009.
- [13] M.Eshaghi Gordji, H.Khodaei and J.M.Rassias, *Fixed point methods for the stability of general quadratic functional equation*, Fixed Point Theory, 12(1)(2011), 71-82.
- [14] D.H.Hyers, *On the stability of the linear functional equation*, Proc.Nat. Acad.Sci.,U.S.A., 27(1941), 222-224.
- [15] D.H.Hyers, G.Isac and Th.M.Rassias, *Stability of functional equations in several variables*, Birkhauser, Basel, (1998).
- [16] Sun Sook Jin and Yang-Hi Lee, *A Fixed Point Approach to the Stability of the Cauchy Additive and Quadratic Type Functional Equation*, Journal of Applied Mathematics, doi:10.1155/2011/817079, 16 pages.
- [17] Sun Sook Jin and Yang Hi Lee, *Fuzzy Stability of a Quadratic-Additive Functional Equation*, International Journal of Mathematics and Mathematical Sciences, doi:10.1155/2011/504802, 16 pages

- [18] K.W.Jun and H.M.Kim, *On the Hyers-Ulam-Rassias stability of a generalized quadratic and additive type functional equation*, Bull. Korean Math. Soc., 42(1)(2005), 133-148.
- [19] K.W.Jun and H.M.Kim, *On the stability of an  $n$ -dimensional quadratic and additive type functional equation*, Math. Ineq. Appl., 9(1)(2006), 153-165.
- [20] S.M. Jung, *On the Hyers-Ulam stability of the functional equations that have the quadratic property*, J. Math. Anal. Appl. 222(1998), 126-137.
- [21] S.M.Jung, *Hyers-Ulam-Rassias Stability of Functional Equations in Mathematical Analysis*, Hadronic Press, Palm Harbor, (2001).
- [22] P.I.Kannappan, *Quadratic functional equation inner product spaces*, Results Math., 27(3-4)(1995), 368-372.
- [23] P.I.Kannappan, *Functional Equations and Inequalities with Applications*, Springer Monographs in Mathematics, (2009).
- [24] B.Margolis and J.B.Diaz, *A fixed point theorem of the alternative for contractions on a generalized complete metric space*, Bull.Amer. Math. Soc., 126(74)(1968), 305-309.
- [25] S.Murthy, M.Arunkumar, G.Ganapathy and P.Rajarethinam, *Stability of mixed type additive quadratic functional equation in Random Normed space*, International Journal of Applied Mathematics, 26(2)(2013), 123-136.
- [26] A.Najati and M.B.Moghimi, *On the stability of a quadratic and additive functional equation*, J. Math. Anal. Appl., 337(2008), 399-415.
- [27] Choonkil Park, *Orthogonal Stability of an Additive-Quadratic Functional Equation*, Fixed Point Theory and Applications 2011(2011), 66.
- [28] Matina J.Rassias, M.Arunkumar and S.Ramamoorthi, *Stability of the Leibniz additive-quadratic functional equation in Quasi-Beta normed space: Direct and fixed point methods*, Journal of Concrete And Applicable Mathematics, 14(1-2)(2014).
- [29] J.M.Rassias, *On approximately of approximately linear mappings by linear mappings*, J. Funct. Anal. USA, 46(1982), 126-130.
- [30] J.M.Rassias, *On approximately of approximately linear mappings by linear mappings*, Bull. Sc. Math., 108(1984), 445-446.
- [31] Th.M.Rassias, *On the stability of the linear mapping in Banach spaces*, Proc.Amer.Math. Soc., 72(1978), 297-300.
- [32] Th.M.Rassias, *Functional Equations, Inequalities and Applications*, Kluwer Academic Publishers, Dordrecht, Bostan London, (2003).
- [33] K.Ravi, M.Arunkumar and J.M.Rassias, *On the Ulam stability for the orthogonally general Euler-Lagrange type functional equation*, International Journal of Mathematical Sciences, 3(08)(2008), 36-47.
- [34] S.M.Ulam, *Problems in Modern Mathematics*, Science Editions, Wiley, NewYork, (1964).
- [35] G.Zamani Eskandani, Hamid Vaezi and Y.N.Dehghan, *Stability Of A Mixed Additive And Quadratic Functional Equation In Non-Archimedean Banach Modules*, Taiwanese Journal Of Mathematics, 14(4)(2010), 1309-1324.