



# Two Hypergeometric Generating Relations Via Gould’s Identity and Their Generalizations

M. I. Qureshi<sup>1</sup> and Sulakshana Bajaj<sup>2,\*</sup>

- 1 Department of Applied Sciences and Humanities, Faculty of Engineering and Technology, Jamia Millia Islamia, New Delhi, India.
- 2 Department of Applied Sciences and Humanities, Modi Institute of Technology, Rawatbhata Road, Kota, Rajasthan, India.

**Abstract:** In the present paper, we have obtained hypergeometric generating relations associated with two hypergeometric polynomials of one variable  $H_n^{(\alpha,\beta)}(x; m)$  and  $\mathcal{B}_n^{(\alpha,\beta)}(x; m, \lambda, \mu)$  with their independent demonstrations via Gould’s identity. As applications, some well known and new generating relations are deduced. Using bounded sequences, further generalizations of two main hypergeometric generating relations have also been given for two generalized polynomials  $S_n^{(\alpha,\beta)}(x; m)$  and  $T_n^{(\alpha,\beta)}(x; m, \lambda, \mu)$ .

**MSC:** 33C05, 33C45, 33C65.

**Keywords:** Jacobi Polynomials, generalized Laguerre polynomial, generalized Rice polynomial of Khandekar, Gould’s identity.  
© JS Publication.

## 1. Introduction and Preliminaries

Throughout in the present paper, we use the following standard notations:  $\mathbb{N} := \{1, 2, 3, \dots\}$ ,  $\mathbb{N}_0 := \{0, 1, 2, 3, \dots\} = \mathbb{N} \cup \{0\}$ ,  $\mathbb{Z}_0^- := \{0, -1, -2, -3, \dots\}$ ,  $\mathbb{Z}^- := \{-1, -2, -3, \dots\} = \mathbb{Z}_0^- \setminus \{0\}$  and  $\mathbb{Z} = (\mathbb{Z}_0^- \cup \mathbb{N})$ . Here, as usual,  $\mathbb{Z}$  denotes the set of integers,  $\mathbb{R}$  denotes the set of real numbers,  $\mathbb{R}^+$  denotes the set of positive real numbers and  $\mathbb{C}$  denotes the set of complex numbers. The Pochhammer symbol (or the shifted factorial)  $(\lambda)_\nu$  ( $\lambda, \nu \in \mathbb{C}$ ) is defined, in terms of the familiar Gamma function, by

$$(\lambda)_\nu := \frac{\Gamma(\lambda + \nu)}{\Gamma(\lambda)} = \begin{cases} 1 & (\nu = 0; \lambda \in \mathbb{C} \setminus \{0\}) \\ \lambda(\lambda + 1) \dots (\lambda + \nu - 1) & (\nu = n \in \mathbb{N}; \lambda \in \mathbb{C}) \end{cases} \tag{1}$$

it is being understood *conventionally* that  $(0)_0 = 1$  and assumed tacitly that the Gamma quotient exists. Some useful consequences of Lagrange’s expansion [21] include the following generalization [10] of the familiar binomial expansion :

$$\sum_{n=0}^{\infty} \binom{\theta + (\beta + 1)n}{n} t^n = \frac{(1 + \zeta)^{\theta + 1}}{(1 - \beta\zeta)} \tag{2}$$

where  $\binom{\theta + (\beta + 1)n}{n}$  is a binomial coefficient and  $\theta, \beta$  are complex numbers independent of  $n$  and  $\zeta$  is a function of ‘t’ defined implicitly by

$$\zeta = t(1 + \zeta)^{1 + \beta} \tag{3}$$

\* E-mail: [sulakshana80@gmail.com](mailto:sulakshana80@gmail.com)

subject to the condition

$$\zeta(0) = 0 \quad (4)$$

Another generalization [10] related with the equation (2), is given as:

$$\sum_{n=0}^{\infty} \frac{\theta}{\{\theta + (\beta + 1)n\}} \binom{\theta + (\beta + 1)n}{n} t^n = 1 + \theta \sum_{n=1}^{\infty} \binom{\theta + (\beta + 1)n - 1}{n - 1} \frac{t^n}{n} = (1 + \zeta)^\theta \quad (5)$$

where  $\zeta$  is defined by the equations (3) and (4). When  $\beta = -1$ , both results (2) and (5) reduce immediately to the binomial expansion. Gould [6] gave the following identity:

$$\sum_{n=0}^{\infty} \frac{\theta(\sigma + \mu n)}{\{\theta + (\beta + 1)n\}} \binom{\theta + (\beta + 1)n}{n} t^n = (1 + \zeta)^\theta \left( \sigma + \frac{\mu\theta\zeta}{1 - \beta\zeta} \right) \quad (6)$$

where  $\theta, \beta, \sigma, \mu$  are complex parameters independent of  $n$  and  $\zeta$  is given by the equations (3) and (4). If we put  $\theta = \{\alpha + (\beta + 1)mr\}$  and  $\sigma = \{\lambda + \mu mr\}$  in Gould's identity (6), we get the first modified form of Gould's identity:

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\{\alpha + (\beta + 1)mr\}(\lambda + \mu n + \mu mr)}{\{\alpha + (\beta + 1)mr + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)mr + (\beta + 1)n}{n} t^n \\ = (1 + \zeta)^{\{\alpha + (\beta + 1)mr\}} \left( \lambda + \mu mr + \frac{\mu\{\alpha + (\beta + 1)mr\}\zeta}{(1 - \beta\zeta)} \right) \end{aligned} \quad (7)$$

with  $\zeta = t(1 + \zeta)^{\beta+1}$ ;  $\zeta(0) = 0$ . If we put  $\theta = \{\alpha + (\beta + 1)mr\}$  and  $\sigma = \{\lambda + \mu(\beta + 1)r\}$  in Gould's identity (6), we get the Second modified form of Gould's identity:

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\{\alpha + (\beta + 1)mr\}\{\lambda + \mu n + \mu(\beta + 1)r\}}{\{\alpha + (\beta + 1)mr + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)mr + (\beta + 1)n}{n} t^n \\ = (1 + \zeta)^{\{\alpha + (\beta + 1)mr\}} \left( \lambda + \mu(\beta + 1)r + \frac{\mu\{\alpha + (\beta + 1)mr\}\zeta}{(1 - \beta\zeta)} \right) \end{aligned} \quad (8)$$

with  $\zeta = t(1 + \zeta)^{\beta+1}$ ;  $\zeta(0) = 0$ .

Gauss's Multiplication Theorem: For every positive integer  $m$ , we have

$$(b)_{mr} = m^{mr} \prod_{j=1}^m \binom{b+j-1}{m}_r \quad ; \quad r = 0, 1, 2, \dots \quad (9)$$

Summation identity [20]

$$\sum_{n=0}^{\infty} \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} B(r, n) = \sum_{n=0}^{\infty} \sum_{r=0}^{\infty} B(r, n + mr) \quad (10)$$

( $\lfloor x \rfloor$  denotes the greatest integer in  $x$ ;  $m \in \mathbb{N}$ ), provided that series involved are absolutely convergent. The generalized Laguerre polynomials  $L_n^{(\alpha)}(x)$  are defined by

$$L_n^{(\alpha)}(x) = \frac{(1 + \alpha)_n}{n!} {}_1F_1 \left[ -n; 1 + \alpha; x \right]; \quad n \in N_0 \quad (11)$$

Replacing  $\alpha$  by  $\alpha + n\beta$  in equation (11), we get

$${}_1F_1 \left[ \begin{matrix} -n & ; \\ 1 + \alpha + n\beta & ; \end{matrix} x \right] = \frac{n!}{(1 + \alpha + n\beta)_n} L_n^{(\alpha+n\beta)}(x) \quad (12)$$

The Jacobi Polynomials of first kind  $P_n^{(\alpha,\beta)}(x)$  [11] are defined by the following equations:

$$P_n^{(\alpha,\beta)}(x) = \frac{(1+\alpha)_n}{n!} {}_2F_1 \left[ \begin{matrix} -n, 1+\alpha+\beta+n; \\ 1+\alpha \end{matrix}; \frac{1-x}{2} \right] \tag{13}$$

$$P_n^{(\alpha,\beta)}(x) = \frac{(1+\alpha+\beta)_{2n}}{n!(1+\alpha+\beta)_n} \left(\frac{x-1}{2}\right)^n {}_2F_1 \left[ \begin{matrix} -n, -\alpha-n; \\ -\alpha-\beta-2n; \end{matrix}; \frac{2}{1-x} \right] \tag{14}$$

where  $n$  is a non-negative integer. Replacing  $\alpha$  by  $(\alpha + bn)$  and  $\beta$  by  $\{\beta - (b + 1)n\}$  in equation (13), we get

$${}_2F_1 \left[ \begin{matrix} -n, 1+\alpha+\beta; \\ 1+\alpha+bn; \end{matrix}; \frac{1-x}{2} \right] = \frac{n! \Gamma(\alpha + bn + 1)}{\Gamma\{\alpha + (b + 1)n + 1\}} P_n^{(\alpha+bn, \beta-(b+1)n)}(x) \tag{15}$$

Replacing  $\alpha$  by  $(\alpha - n)$  and  $\beta$  by  $\{\beta - (b + 1)n\}$  in equation (14), we get the following result

$${}_2F_1 \left[ \begin{matrix} -n, -\alpha; \\ -\alpha-\beta+bn; \end{matrix}; \frac{2}{1-x} \right] = \frac{\Gamma(1+\alpha+\beta-bn-n)n!}{\Gamma(1+\alpha+\beta-bn)} \left(\frac{2}{x-1}\right)^n P_n^{(\alpha-n, \beta-bn-n)}(x) \tag{16}$$

The generalized Rice Polynomials  $H_n^{(\alpha,\beta)}[\nu, \sigma, x]$  of Khandekar [9] are defined by

$$H_n^{(\alpha,\beta)}[\nu, \sigma, x] = \binom{\alpha+n}{n} {}_3F_2 \left[ \begin{matrix} -n, \alpha+\beta+n+1, \nu; \\ \alpha+1, \sigma \end{matrix}; x \right] \tag{17}$$

$$H_n[\nu, \sigma, x] = H_n^{(0,0)}[\nu, \sigma, x] \tag{18}$$

$$P_n^{(\alpha,\beta)}(x) = H_n^{(\alpha,\beta)} \left[ \nu, \nu, \frac{1-x}{2} \right] \tag{19}$$

Replacing  $\alpha$  by  $(\alpha + bn)$  and  $\beta$  by  $\{\beta - (b + 1)n\}$  in Equation (17), we get

$${}_3F_2 \left[ \begin{matrix} -n, \nu, 1+\alpha+\beta; \\ 1+\alpha+bn, \sigma; \end{matrix}; x \right] = \frac{n!}{(1+\alpha+bn)_n} H_n^{(\alpha+bn, \beta-(b+1)n)}[\nu, \sigma, x] \tag{20}$$

Some useful Pochhammer’s relations

$$\frac{(\lambda + \mu n) \left(\frac{\lambda+\mu n}{\mu(\beta+1-m)} + 1\right)_r}{\left(\frac{\lambda+\mu n}{\mu(\beta+1-m)}\right)_r} = \lambda + \mu n + \mu(\beta + 1)r - \mu m r \tag{21}$$

$$\mu(\beta + 1)r + \frac{\mu\zeta}{(1 - \beta\zeta)} \{\alpha + m(\beta + 1)r\} = \frac{\mu\zeta\alpha}{(1 - \beta\zeta)} \frac{\left(\frac{\zeta\alpha}{(\beta+1)(1+m\zeta-\beta\zeta)} + 1\right)_r}{\left(\frac{\zeta\alpha}{(\beta+1)(1+m\zeta-\beta\zeta)}\right)_r} \tag{22}$$

$$\alpha + m(\beta + 1)r = \alpha \frac{\left(\frac{\alpha}{m(\beta+1)} + 1\right)_r}{\left(\frac{\alpha}{m(\beta+1)}\right)_r} \tag{23}$$

$$\mu m r + \frac{\mu\zeta\{\alpha + (\beta + 1)m r\}}{(1 - \beta\zeta)} = \frac{\mu\zeta}{(1 - \beta\zeta)} \frac{\left(\frac{\alpha\zeta}{m(1+\zeta)} + 1\right)_r}{\left(\frac{\alpha\zeta}{m(1+\zeta)}\right)_r} \tag{24}$$

where  $r = 0, 1, 2, 3, \dots$ . Now we shall discuss some special cases of the implicit functions defined by equation (3) subject to the condition (5). Using Mathematica 9.0, we can find the roots of resulting cubic equation in  $\zeta$  for different values of  $\beta$  in

equation (3).

**Case I:-** When  $\beta = 0$  in (3), then particular value of  $\zeta$  (satisfying the condition (4)) is denoted by

$$\Theta = \frac{t}{1-t} \quad (25)$$

**Case II:-** When  $\beta = 1$  in (3), we get  $t\zeta^2 + (2t-1)\zeta + t = 0$ , then one of the values of  $\zeta$  (satisfying the condition (4)) is given by

$$\Lambda = \frac{1-2t-\sqrt{(1-4t)}}{2t} \quad (26)$$

**Case III:-** When  $\beta = -2$  in (3), we get  $\zeta^2 + \zeta - t = 0$ , then the particular value of  $\zeta$  (satisfying the condition (4)) is given by

$$\Xi = \frac{-1+\sqrt{(1+4t)}}{2} \quad (27)$$

**Case IV:-** When  $\beta = -3$  in (3), we get  $\zeta^3 + 2\zeta^2 + \zeta - t = 0$ , then one of the roots (satisfying the condition (4)) of above equation is given by

$$\Upsilon = \frac{1}{3} \left[ -2 + \frac{2^{\frac{1}{3}}}{\left\{2+27t+3\sqrt{3}\sqrt{(4t+27t^2)}\right\}^{\frac{1}{3}}} + \frac{\left\{2+27t+3\sqrt{3}\sqrt{(4t+27t^2)}\right\}^{\frac{1}{3}}}{2^{\frac{1}{3}}} \right] \quad (28)$$

**Case V:-** When  $\beta = -\frac{1}{2}$  in (3), we get  $\zeta^2 - t^2\zeta - t^2 = 0$ , then one of the roots (satisfying the condition (4)) of above equation is given by

$$U = \frac{t}{2} \{t + \sqrt{(t^2+4)}\} \quad (29)$$

**Case VI:-** When  $\beta = \frac{-3}{2}$  in (3), we get  $\zeta^3 + \zeta^2 - t^2 = 0$ , then one of the roots (satisfying the condition (4)) of above equation is given by

$$\Psi = -\frac{1}{3} \left[ 1 + \frac{(1+\iota\sqrt{3})}{2^{\frac{2}{3}} \left\{-2+27t^2+3\sqrt{3}\sqrt{(-4t^2+27t^4)}\right\}^{\frac{1}{3}}} + \frac{(1-\iota\sqrt{3}) \left\{-2+27t^2+3\sqrt{3}\sqrt{(-4t^2+27t^4)}\right\}^{\frac{1}{3}}}{2^{\frac{4}{3}}} \right] \quad (30)$$

where  $\iota = \sqrt{(-1)}$ .

**Case VII:-** When  $\beta = -\frac{1}{3}$  in (3), we obtain  $\zeta^3 - t^3\zeta^2 - 2t^3\zeta - t^3 = 0$ , then one of the values of  $\zeta$  (satisfying the condition (4)) is denoted by

$$\Pi = \frac{t^3}{3} - \frac{2^{\frac{1}{3}}(-6t^3-t^6)}{3\{27t^3+18t^6+2t^9+3\sqrt{3}\sqrt{(27t^6+4t^9)}\}^{\frac{1}{3}}} + \frac{\{27t^3+18t^6+2t^9+3\sqrt{3}\sqrt{(27t^6+4t^9)}\}^{\frac{1}{3}}}{3.2^{\frac{1}{3}}} \quad (31)$$

**Case VIII:-** When  $\beta = -\frac{2}{3}$  in (3), we obtain  $\zeta^3 - t^3\zeta - t^3 = 0$ , then one of the values of  $\zeta$  (satisfying the condition (4)) is denoted by

$$\Phi = \frac{\left(\frac{2}{3}\right)^{\frac{1}{3}} t^3}{\left\{9t^3+\sqrt{3}\sqrt{(27t^6-4t^9)}\right\}^{\frac{1}{3}}} + \frac{\left\{9t^3+\sqrt{3}\sqrt{(27t^6-4t^9)}\right\}^{\frac{1}{3}}}{2^{\frac{1}{3}} 3^{\frac{2}{3}}} \quad (32)$$

## 2. Main Generating Relations

**First Generating Relation:** If any values of variables and parameters leading to the results which do not make sense, are tacitly excluded, then

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{\{\alpha + (\beta + 1)n\}} H_n^{(\alpha, \beta)}(x; m) t^n = (1 + \zeta)^\alpha \left\{ \frac{\lambda}{\alpha} {}_{p+1}F_{q+1} \left[ \begin{matrix} \frac{\alpha}{m(\beta+1)}, (a_p) & ; & x(-\zeta)^m \\ 1 + \frac{\alpha}{m(\beta+1)}, (b_q) & & \end{matrix} \right] \right. \\ \left. + \frac{\mu\zeta}{(1 - \beta\zeta)} {}_{p+2}F_{q+2} \left[ \begin{matrix} \frac{\alpha}{m(\beta+1)}, (a_p), 1 + \frac{\alpha\zeta}{m(\zeta+1)} & ; & x(-\zeta)^m \\ 1 + \frac{\alpha}{m(\beta+1)}, (b_q), \frac{\alpha\zeta}{m(\zeta+1)} & & \end{matrix} \right] \right\} \quad (33)$$

where  $\zeta = t(1 + \zeta)^{\beta+1}$ ;  $\zeta(0) = 0$  provided that involved series on both sides are absolutely convergent. Here Srivastava’s generalized hypergeometric polynomials  $H_n^{(\alpha, \beta)}(x; m)$  [18, 20] are given by

$$H_n^{(\alpha, \beta)}(x; m) = \binom{\alpha + (\beta + 1)n}{n} {}_{p+m}F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) & ; & x \\ \Delta(m; 1 + \alpha + \beta n), (b_q) & & \end{matrix} \right], \quad (34)$$

where  $\alpha$  and  $\beta$  are complex parameters independent of ‘ $n$ ’ and  $\Delta(m; \lambda)$  abbreviates the array of  $m$  number of parameters given by

$$\frac{\lambda}{m}, \frac{\lambda + 1}{m}, \dots, \frac{\lambda + m - 1}{m}; m \in \mathbb{N}$$

**Independent Demonstration:** Using definition (34) of  $H_n^{(\alpha, \beta)}(x; m)$  and then the power series form of  ${}_{p+m}F_{q+m}[x]$  in left hand side of Equation (33), we get

$$\Omega = \sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{\{\alpha + (\beta + 1)n\}} H_n^{(\alpha, \beta)}(x; m) t^n \\ = \sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{\{\alpha + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)n}{n} {}_{p+m}F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p); & x \\ \Delta(m; 1 + \alpha + n\beta), (b_q); & \end{matrix} \right] t^n \\ = \sum_{n=0}^{\infty} \frac{(\lambda + \mu n) \Gamma\{\alpha + (\beta + 1)n + 1\}}{\{\alpha + (\beta + 1)n\} \Gamma(n + 1) \Gamma\{\alpha + \beta n + 1\}} \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} \frac{\prod_{j=1}^m \binom{-n + j - 1}{m} [(a_p)]_r x^r t^n}{\prod_{j=1}^m \binom{1 + \alpha + n\beta + j - 1}{m} [(b_q)]_r r!}$$

Using Gauss’s multiplication theorem (9) in above equation, we get

$$\Omega = \sum_{n=0}^{\infty} \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} \frac{(\lambda + \mu n) (-n)_{mr} (\alpha)_{n(\beta+1)} [(a_p)]_r x^r t^n}{(\alpha + 1)_{n\beta} (1 + \alpha + n\beta)_{mr} [(b_q)]_r r! n!} \quad (35)$$

Now applying summation identity (10) and then simplifying further, we get

$$\Omega = \sum_{r=0}^{\infty} \frac{(\alpha)_{m(\beta+1)r} [(a_p)]_r x^r (-t)^{mr}}{(\alpha + 1)_{m(\beta+1)r} [(b_q)]_r r!} \sum_{n=0}^{\infty} \frac{\{\alpha + (\beta + 1)mr\} (\lambda + \mu n + \mu mr)}{\{\alpha + (\beta + 1)mr + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)mr + (\beta + 1)n}{n} t^n \quad (36)$$

Now using first modified Gould’s identity (7), we get

$$\Omega = (1 + \zeta)^\alpha \sum_{r=0}^{\infty} \frac{(\alpha)_{m(\beta+1)r} [(a_p)]_r x^r (-t)^{mr}}{(\alpha + 1)_{m(\beta+1)r} [(b_q)]_r r!} (1 + \zeta)^{m(\beta+1)r} \cdot \left[ \lambda + \left( \mu m r + \frac{\mu\zeta\{\alpha + (\beta + 1)mr\}}{(1 - \beta\zeta)} \right) \right] \quad (37)$$

Simplifying it further and using equation (3), (4) and (24), we get

$$\Omega = \lambda (1 + \zeta)^\alpha \sum_{r=0}^{\infty} \frac{\left(\frac{\alpha}{m(\beta+1)}\right)_r [(a_p)]_r x^r (-\zeta)^{mr}}{\left(\frac{\alpha}{m(\beta+1)} + 1\right)_r [(b_q)]_r r!} + (1 + \zeta)^\alpha \frac{\mu \zeta \alpha}{(1 - \beta \zeta)} \sum_{r=0}^{\infty} \frac{\left(\frac{\alpha}{m(\beta+1)}\right)_r [(a_p)]_r \left(\frac{\alpha \zeta}{m(\zeta+1)} + 1\right)_r x^r (-\zeta)^{mr}}{\left(\frac{\alpha}{m(\beta+1)} + 1\right)_r [(b_q)]_r \left(\frac{\alpha \zeta}{m(\zeta+1)}\right)_r r!} \quad (38)$$

after solving it further, we get the desired result (33).

**Second Generating Relation:** If any values of variables and parameters leading to the results which do not make sense, are tacitly excluded, then

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{\{\alpha + (\beta + 1)n\}} \mathcal{B}_n^{(\alpha, \beta)}(x; m, \lambda, \mu) t^n = (1 + \zeta)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{\alpha}{m(\beta+1)}, (a_p); \\ 1 + \frac{\alpha}{m(\beta+1)}, (b_q); \end{matrix} x(-\zeta)^m \right] \right. \\ \left. + \frac{\mu \zeta}{(1 - \beta \zeta)} {}_{p+2}F_{q+2} \left[ \begin{matrix} (a_p), \frac{\alpha}{m(\beta+1)}, \frac{\zeta \alpha}{(\beta+1)(1+m\zeta-\beta\zeta)} + 1; \\ (b_q), \frac{\alpha}{m(\beta+1)} + 1, \frac{\zeta \alpha}{(\beta+1)(1+m\zeta-\beta\zeta)}; \end{matrix} x(-\zeta)^m \right] \right\} \quad (39)$$

where  $\zeta = t(1 + \zeta)^{\beta+1}$ ;  $\zeta(0) = 0$  provided that involved series on both sides are absolutely convergent. Here we define new generalized hypergeometric polynomials  $\mathcal{B}_n^{(\alpha, \beta)}(x; m, \lambda, \mu)$ , known as ‘‘Pathan’s generalized hypergeometric polynomials of one variable’’, given by

$$\mathcal{B}_n^{(\alpha, \beta)}(x; m, \lambda, \mu) = \binom{\alpha + (\beta + 1)n}{n} {}_{p+m+1}F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 + \frac{\lambda + \mu n}{\mu(\beta+1-m)}, (a_p); \\ \Delta(m; 1 + \alpha + \beta n), \frac{\lambda + \mu n}{\mu(\beta+1-m)}, (b_q); \end{matrix} x \right] \quad (40)$$

where  $\alpha$  and  $\beta$  are complex parameters independent of ‘ $n$ ’ and  $\Delta(m; \lambda)$  abbreviates the array of  $m$  number of parameters given by

$$\frac{\lambda}{m}, \frac{\lambda + 1}{m}, \dots, \frac{\lambda + m - 1}{m}; m \in \mathbb{N}$$

**Independent Demonstration:** Using the definition (40) of  $\mathcal{B}_n^{(\alpha, \beta)}(x; m, \lambda, \mu)$  and then the power series form of  ${}_{p+m+1}F_{q+m+1}[x]$  in left hand side of equation (39), using Gauss’s multiplication Theorem 1.9 and Result 1.21, we get

$$\Omega_1 = \sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{\{\alpha + (\beta + 1)n\}} \mathcal{B}_n^{(\alpha, \beta)}(x; m, \lambda, \mu) t^n \\ = \sum_{n=0}^{\infty} \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} \frac{[\lambda + \mu n + \mu(\beta + 1)r - \mu m r] (-n)_{mr} \Gamma\{\alpha + n(\beta + 1)\} [(a_p)]_r x^r t^n}{\Gamma\{\alpha + n\beta + 1 + mr\} [(b_q)]_r r! n!} \quad (41)$$

Now applying summation identity (10) in above equation then simplifying further, we get

$$\Omega_1 = \sum_{r=0}^{\infty} \frac{[(a_p)]_r x^r (-t)^{mr}}{[\alpha + m(\beta + 1)r] [(b_q)]_r r!} \sum_{n=0}^{\infty} \frac{\{\alpha + (\beta + 1)mr\} (\lambda + \mu n + \mu(\beta + 1)r)}{\{\alpha + (\beta + 1)mr + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)mr + (\beta + 1)n}{n} t^n \quad (42)$$

Now using second modified Gould’s identity (8), we get

$$\Omega_1 = (1 + \zeta)^\alpha \sum_{r=0}^{\infty} \frac{[(a_p)]_r x^r (-t)^{mr} (1 + \zeta)^{m(\beta+1)r}}{\{\alpha + m(\beta + 1)r\} [(b_q)]_r r!} \left[ \lambda + \left( \mu(\beta + 1)r + \frac{\mu \zeta}{(1 - \beta \zeta)} \{\alpha + m(\beta + 1)r\} \right) \right] \quad (43)$$

Now using equation (22) and (23) in above equation and summing it up into hypergeometric form further, we get the desired result (39).

### 3. Known Applications of Generating Relation (33)

(i). Putting  $\lambda = 1, \mu = 0$  in equation (33), we get

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\alpha}{\{\alpha + (\beta + 1)n\}} H_n^{(\alpha, \beta)}(x; m) t^n &= \sum_{n=0}^{\infty} \frac{\alpha}{\{\alpha + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)n}{n} {}_{p+m}F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) & ; & x \\ \Delta(m; 1 + \alpha + n\beta), (b_q) & & \end{matrix} \right] t^n \\ &= (1 + \zeta)^\alpha {}_{p+1}F_{q+1} \left[ \begin{matrix} \frac{\alpha}{m(\beta+1)}, (a_p); & & x(-\zeta)^m \\ 1 + \frac{\alpha}{m(\beta+1)}, (b_q); & & \end{matrix} \right] \end{aligned} \tag{44}$$

with  $\zeta = t(1 + \zeta)^{\beta+1}; \zeta(0) = 0$  which is the more simplified form of a result of Srivastava [20].

(ii). Putting  $\lambda = 1, \mu = 0$  and  $m = 1$  in equation (2.1), we get

$$\sum_{n=0}^{\infty} \frac{\alpha}{\{\alpha + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)n}{n} {}_{p+1}F_{q+1} \left[ \begin{matrix} -n, (a_p); & & x \\ 1 + \alpha + n\beta, (b_q); & & \end{matrix} \right] t^n = (1 + \zeta)^\alpha {}_{p+1}F_{q+1} \left[ \begin{matrix} \frac{\alpha}{(\beta+1)}, (a_p); & & -x\zeta \\ 1 + \frac{\alpha}{(\beta+1)}, (b_q); & & \end{matrix} \right] \tag{45}$$

with  $\zeta = t(1 + \zeta)^{\beta+1}; \zeta(0) = 0$  which is the result of Brown [3].

(iii). Putting  $\lambda = 1, \mu = \frac{\beta+1}{\alpha}$  in equation (33), we get

$$\begin{aligned} \sum_{n=0}^{\infty} \binom{\alpha + (\beta + 1)n}{n} {}_{p+m}F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) & ; & x \\ \Delta(m; 1 + \alpha + n\beta), (b_q); & & \end{matrix} \right] t^n &= (1 + \zeta)^\alpha {}_{p+1}F_{q+1} \left[ \begin{matrix} \frac{\alpha}{m(\beta+1)}, (a_p) & ; & x(-\zeta)^m \\ 1 + \frac{\alpha}{m(\beta+1)}, (b_q); & & \end{matrix} \right] \\ &+ \frac{(\beta + 1)\zeta(1 + \zeta)^\alpha}{(1 - \beta\zeta)} {}_{p+2}F_{q+2} \left[ \begin{matrix} \frac{\alpha}{m(\beta+1)}, (a_p), 1 + \frac{\alpha\zeta}{m(\zeta+1)}; & & x(-\zeta)^m \\ 1 + \frac{\alpha}{m(\beta+1)}, (b_q), \frac{\alpha\zeta}{m(\zeta+1)}; & & \end{matrix} \right] \end{aligned}$$

After further simplification, we have

$$\sum_{n=0}^{\infty} H_n^{(\alpha, \beta)}(x; m) t^n = \frac{(1 + \zeta)^{(\alpha+1)}}{(1 - \beta\zeta)} {}_pF_q \left[ \begin{matrix} (a_p) & ; & x(-\zeta)^m \\ (b_q) & ; & \end{matrix} \right] \tag{46}$$

with  $\zeta = t(1 + \zeta)^{\beta+1}; \zeta(0) = 0$ , which is the known result of Srivastava [16, 17].

(iv). Putting  $\lambda = 1, \mu = \frac{\beta+1}{\alpha}$  and  $m = 1$  in equation (2.1), we get

$$\begin{aligned} \sum_{n=0}^{\infty} \binom{\alpha + (\beta + 1)n}{n} {}_{p+1}F_{q+1} \left[ \begin{matrix} -n, (a_p) & ; & x \\ 1 + \alpha + n\beta, (b_q); & & \end{matrix} \right] t^n &= (1 + \zeta)^\alpha {}_{p+1}F_{q+1} \left[ \begin{matrix} \frac{\alpha}{(\beta+1)}, (a_p); & & -x\zeta \\ 1 + \frac{\alpha}{(\beta+1)}, (b_q); & & \end{matrix} \right] \\ &+ \frac{(\beta + 1)\zeta(1 + \zeta)^\alpha}{(1 - \beta\zeta)} {}_{p+2}F_{q+2} \left[ \begin{matrix} \frac{\alpha}{(\beta+1)}, (a_p), 1 + \frac{\alpha\zeta}{(\zeta+1)}; & & -x\zeta \\ 1 + \frac{\alpha}{(\beta+1)}, (b_q), \frac{\alpha\zeta}{(\zeta+1)}; & & \end{matrix} \right] \\ &= \frac{(1 + \zeta)^{(\alpha+1)}}{(1 - \beta\zeta)} {}_pF_q \left[ \begin{matrix} (a_p) & ; & -x\zeta \\ (b_q) & ; & \end{matrix} \right] \end{aligned} \tag{47}$$

with  $\zeta = t(1 + \zeta)^{\beta+1}; \zeta(0) = 0$  which is the known result of Srivastava [14, 15].

(v). Putting  $\lambda = 1, \mu = 0, \beta = \frac{-1}{2}, m = 1$  and replacing  $\zeta$  by  $U$  in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{\alpha}{(\alpha + \frac{n}{2})} f_n^{(\alpha)}(x) t^n = \sum_{n=0}^{\infty} \frac{\alpha}{(\alpha + \frac{n}{2})} \binom{\alpha + \frac{n}{2}}{n} {}_{p+1}F_{q+1} \left[ \begin{matrix} -n, (a_p) & ; & x \\ 1 + \alpha - \frac{n}{2}, (b_q); & & \end{matrix} \right] t^n$$

$$= (1 + U)^\alpha {}_{p+1}F_{q+1} \left[ \begin{matrix} 2\alpha, (a_p); \\ 1 + 2\alpha, (b_q); \end{matrix} -xU \right] \tag{48}$$

where  $U = \frac{t}{2}\{t + \sqrt{t^2 + 4}\}$ . This is well known Brown's result [2].

(vi). Putting  $\lambda = 1, \mu = \frac{1}{2\alpha}, \beta = \frac{-1}{2}, m = 1$  and replacing  $\zeta$  by  $U$  in equation (33), we get

$$\begin{aligned} \sum_{n=0}^\infty f_n^{(\alpha)}(x)t^n &= \sum_{n=0}^\infty \binom{\alpha + \frac{n}{2}}{n} {}_{p+1}F_{q+1} \left[ \begin{matrix} -n, (a_p) & ; & x \\ 1 + \alpha - \frac{n}{2}, (b_q); \end{matrix} \right] t^n \\ &= (1 + U)^\alpha {}_{p+1}F_{q+1} \left[ \begin{matrix} 2\alpha, (a_p); \\ 1 + 2\alpha, (b_q); \end{matrix} -xU \right] + \frac{U(1 + U)^\alpha}{2(1 + \frac{1}{2}U)} {}_{p+2}F_{q+2} \left[ \begin{matrix} 2\alpha, (a_p), 1 + \frac{\alpha U}{1+U}; \\ 1 + 2\alpha, (b_q), \frac{\alpha U}{(U+1)}; \end{matrix} -xU \right] \\ &= \frac{(1 + U)^{\alpha+1}}{(1 + \frac{1}{2}U)^p} {}_pF_q \left[ \begin{matrix} (a_p); \\ (b_q); \end{matrix} -xU \right] \end{aligned} \tag{49}$$

where  $U = \frac{t}{2}\{t + \sqrt{t^2 + 4}\}$ . It is also a well known result of Brown [2].

### 4. Some Special Cases of Generating Relation (45)

(i). Taking  $p = 0 = q$  in (45), we get

$$\sum_{n=0}^\infty \frac{\alpha}{\{\alpha + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)n}{n} {}_1F_1 \left[ \begin{matrix} -n & ; & x \\ 1 + \alpha + n\beta; \end{matrix} \right] t^n = (1 + \zeta)^\alpha {}_1F_1 \left[ \begin{matrix} \frac{\alpha}{(\beta+1)}; \\ 1 + \frac{\alpha}{(\beta+1)}; \end{matrix} -x\zeta \right] \tag{50}$$

Now using the definition of generalized Laguerre polynomial (12) and solving we get

$$\sum_{n=0}^\infty \frac{\alpha}{\{\alpha + (\beta + 1)n\}} L_n^{(\alpha+\beta n)}(x)t^n = (1 + \zeta)^\alpha {}_1F_1 \left[ \begin{matrix} \frac{\alpha}{(\beta+1)} & ; & -x\zeta \\ 1 + \frac{\alpha}{(\beta+1)}; \end{matrix} \right] \tag{51}$$

In equations (50) and (51),  $\zeta$  is given by the equations (3), (4). It is the known generating relation of Brown [4, 13, 19].

(ii). In equation (45), putting  $p = 1, q = 0$ , replacing  $x$  by  $\frac{1-x}{2}, \beta$  by  $b$  and  $\zeta$  by  $\xi$ , we get

$$\sum_{n=0}^\infty \frac{\alpha}{\{\alpha + (b + 1)n\}} \binom{\alpha + (b + 1)n}{n} {}_2F_1 \left[ \begin{matrix} -n, a_1 & ; & \frac{(1-x)}{2} \\ 1 + \alpha + bn; \end{matrix} \right] t^n = (1 + \xi)^\alpha {}_2F_1 \left[ \begin{matrix} \frac{\alpha}{b+1}, a_1 & ; & \xi \frac{(x-1)}{2} \\ 1 + \frac{\alpha}{b+1}; \end{matrix} \right] \tag{52}$$

where  $\xi$  is a function of 't', defined implicitly by

$$\xi = t(1 + \xi)^{b+1} \quad ; \xi(0) = 0 \tag{53}$$

Putting  $a_1 = 1 + \alpha + \beta$  in above equation and using equation (15), we get

$$\sum_{n=0}^\infty \frac{\alpha}{\{\alpha + (b + 1)n\}} P_n^{(\alpha+bn, \beta-(b+1)n)}(x)t^n = (1 + \xi)^\alpha {}_2F_1 \left[ \begin{matrix} \frac{\alpha}{b+1}, 1 + \alpha + \beta & ; & \frac{(x-1)\xi}{2} \\ 1 + \frac{\alpha}{b+1}; \end{matrix} \right] \tag{54}$$

where  $\xi$  is given by equation (4.4).

(iii). In equation (45), putting  $p = 1, q = 0, \alpha = a, \beta = b$ , replacing  $x$  by  $\frac{2}{1-x}$  and  $\zeta$  by  $\xi$ , we get

$$\sum_{n=0}^\infty \frac{a}{\{a + (b + 1)n\}} \binom{a + (b + 1)n}{n} {}_2F_1 \left[ \begin{matrix} -n, a_1 & ; & \frac{2}{1-x} \\ 1 + a + bn; \end{matrix} \right] t^n = (1 + \xi)^a {}_2F_1 \left[ \begin{matrix} \frac{a}{b+1}, a_1 & ; & \frac{2\xi}{(x-1)} \\ 1 + \frac{a}{b+1}; \end{matrix} \right] \tag{55}$$



where  $\xi = t(1 + \xi)^{(b+1)}$ ;  $\xi(0) = 0$ . Replacing  $a_1$  by  $-\alpha$  and then  $a$  by  $-\alpha - \beta - 1$ , and using equation (16), we get the following result

$$\sum_{n=0}^{\infty} \frac{(\alpha + \beta + 1)}{\{\alpha + \beta + 1 - (b + 1)n\}} \left(\frac{2}{1-x}\right)^n P_n^{(\alpha-n, \beta-bn-n)}(x) t^n = (1 + \xi)^{-(\alpha+\beta+1)} {}_2F_1 \left[ \begin{matrix} -\frac{(\alpha+\beta+1)}{b+1}, -\alpha; \\ 1 - \frac{(\alpha+\beta+1)}{b+1}; \end{matrix} \frac{2\xi}{(x-1)} \right] \tag{56}$$

where  $\xi = t(1 + \xi)^{(b+1)}$ ;  $\xi(0) = 0$ . Now replacing  $t$  by  $t\frac{(1-x)}{2}$  and  $\xi$  by  $w$ , we get

$$\sum_{n=0}^{\infty} \frac{(\alpha + \beta + 1)}{\{\alpha + \beta + 1 - (b + 1)n\}} P_n^{(\alpha-n, \beta-bn-n)}(x) t^n = (1 + w)^{-(\alpha+\beta+1)} {}_2F_1 \left[ \begin{matrix} -\frac{(\alpha+\beta+1)}{b+1}, -\alpha; \\ 1 - \frac{(\alpha+\beta+1)}{b+1}; \end{matrix} \frac{2w}{(x-1)} \right] \tag{57}$$

where  $w(x, t) = t\frac{(1-x)}{2}(1 + w)^{b+1}$ ;  $w(x, 0) = 0$ .

(iv). putting  $p = 2, q = 1, \beta = b, a_1 = 1 + \alpha + \beta, a_2 = \nu$  and  $b_1 = \sigma$  and replacing  $\zeta$  by  $\xi$  in equation (45), we get

$$\sum_{n=0}^{\infty} \frac{\alpha}{\{\alpha + (b + 1)n\}} \binom{\alpha + (b + 1)n}{n} {}_3F_2 \left[ \begin{matrix} -n, 1 + \alpha + \beta, \nu; \\ 1 + \alpha + n, b, \sigma \end{matrix}; x \right] t^n = (1 + \xi)^\alpha {}_3F_2 \left[ \begin{matrix} \frac{\alpha}{b+1}, 1 + \alpha + \beta, \nu; \\ 1 + \frac{\alpha}{b+1}, \sigma \end{matrix}; -x\xi \right] \tag{58}$$

where  $\xi$  is given by equation (53). Now using equation (20) in above equation, we get:

$$\sum_{n=0}^{\infty} \frac{\alpha}{\{\alpha + (b + 1)n\}} H_n^{(\alpha+bn, \beta-bn-n)}[\nu, \sigma, x] t^n = (1 + \xi)^\alpha {}_3F_2 \left[ \begin{matrix} \frac{\alpha}{b+1}, 1 + \alpha + \beta, \nu; \\ 1 + \frac{\alpha}{b+1}, \sigma \end{matrix}; x(-\xi) \right] \tag{59}$$

where  $\xi$  is given by equation (53).

### 5. New Applications of First Generating Relation (33)

(i). Putting  $\beta = 0$  and  $\zeta = \Theta = \frac{t}{1-t}$  from equation (25) in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + n)} \binom{\alpha + n}{n} {}_{p+m}F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) \\ \Delta(m; 1 + \alpha), (b_q) \end{matrix}; x \right] t^n = (1 - t)^{-\alpha} \left\{ \frac{\lambda}{\alpha} {}_{p+1}F_{q+1} \left[ \begin{matrix} \frac{\alpha}{m}, (a_p); \\ 1 + \frac{\alpha}{m}, (b_q); \end{matrix} x \left(\frac{-t}{1-t}\right)^m \right] + \frac{\mu t}{(1-t)} {}_{p+2}F_{q+2} \left[ \begin{matrix} \frac{\alpha}{m}, 1 + \frac{t\alpha}{m}, (a_p); \\ 1 + \frac{\alpha}{m}, \frac{t\alpha}{m}, (b_q); \end{matrix} x \left(\frac{-t}{1-t}\right)^m \right] \right\} \tag{60}$$

(ii). Putting  $\beta = 1$  and  $\zeta = \Lambda = \frac{1-2t-\sqrt{(1-4t)}}{2t}$  from equation (26) in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + 2n)} \binom{\alpha + 2n}{n} {}_{p+m}F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) \\ \Delta(m; 1 + \alpha + n), (b_q) \end{matrix}; x \right] t^n = (1 + \Lambda)^\alpha \left\{ \frac{\lambda}{\alpha} {}_{p+1}F_{q+1} \left[ \begin{matrix} \frac{\alpha}{2m}, (a_p) \\ 1 + \frac{\alpha}{2m}, (b_q); \end{matrix} x(-\Lambda)^m \right] + \frac{\mu\Lambda}{(1-\Lambda)} {}_{p+2}F_{q+2} \left[ \begin{matrix} \frac{\alpha}{2m}, 1 + \frac{\alpha\Lambda}{m(\Lambda+1)}, (a_p); \\ 1 + \frac{\alpha}{2m}, \frac{\alpha\Lambda}{m(\Lambda+1)}, (b_q); \end{matrix} x(-\Lambda)^m \right] \right\} \tag{61}$$

(iii). Putting  $\beta = -2$  and  $\zeta = \Xi = \frac{-1+\sqrt{(1+4t)}}{2}$  from equation (27) in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha - n)} \binom{\alpha - n}{n} {}_{p+m}F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) \\ \Delta(m; 1 + \alpha - 2n), (b_q) \end{matrix}; x \right] t^n = (1 + \Xi)^\alpha \left\{ \frac{\lambda}{\alpha} {}_{p+1}F_{q+1} \left[ \begin{matrix} \frac{-\alpha}{m}, (a_p); \\ 1 - \frac{\alpha}{m}, (b_q); \end{matrix} x(-\Xi)^m \right] \right\}$$

$$+ \frac{\mu \Xi}{(1+2\Xi)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{-\alpha}{m}, 1 + \frac{\alpha \Xi}{m(\Xi+1)}, (a_p); \\ 1 - \frac{\alpha}{m}, \frac{\alpha \Xi}{m(\Xi+1)}, (b_q); \end{matrix} x(-\Xi)^m \right] \quad (62)$$

(iv). Putting  $\beta = -3$  and  $\zeta = \Upsilon$  from equation (28) in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha - 2n)} \binom{\alpha - 2n}{n}_{p+m} F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) & ; & x \\ \Delta(m; 1 + \alpha - 3n), (b_q); \end{matrix} \right] t^n = (1 + \Upsilon)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{-\alpha}{2m}, (a_p); \\ 1 - \frac{\alpha}{2m}, (b_q); \end{matrix} x(-\Upsilon)^m \right] \right. \\ \left. + \frac{\mu \Upsilon}{(1+3\Upsilon)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{-\alpha}{2m}, 1 + \frac{\alpha \Upsilon}{m(\Upsilon+1)}, (a_p); \\ 1 - \frac{\alpha}{2m}, \frac{\alpha \Upsilon}{m(\Upsilon+1)}, (b_q); \end{matrix} x(-\Upsilon)^m \right] \right\} \quad (63)$$

(v). Putting  $\beta = \frac{-1}{2}$  and  $\zeta = U$  from equation (29) in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + \frac{1}{2}n)} \binom{\alpha + \frac{1}{2}n}{n}_{p+m} F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) & ; & x \\ \Delta(m; 1 + \alpha - \frac{1}{2}n), (b_q); \end{matrix} \right] t^n = (1 + U)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{2\alpha}{m}, (a_p); \\ 1 + \frac{2\alpha}{m}, (b_q); \end{matrix} x(-U)^m \right] \right. \\ \left. + \frac{\mu U}{(1 + \frac{1}{2}U)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{2\alpha}{m}, 1 + \frac{\alpha U}{m(U+1)}, (a_p); \\ 1 + \frac{2\alpha}{m}, \frac{\alpha U}{m(U+1)}, (b_q); \end{matrix} x(-U)^m \right] \right\} \quad (64)$$

(vi). Putting  $\beta = \frac{-3}{2}$  and  $\zeta = \Psi$  from equation (30) in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha - \frac{1}{2}n)} \binom{\alpha - \frac{1}{2}n}{n}_{p+m} F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) & ; & x \\ \Delta(m; 1 + \alpha - \frac{3}{2}n), (b_q); \end{matrix} \right] t^n = (1 + \Psi)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{-2\alpha}{m}, (a_p); \\ 1 - \frac{2\alpha}{m}, (b_q); \end{matrix} x(-\Psi)^m \right] \right. \\ \left. + \frac{\mu \Psi}{(1 + \frac{3}{2}\Psi)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{-2\alpha}{m}, 1 + \frac{\alpha \Psi}{m(\Psi+1)}, (a_p); \\ 1 - \frac{2\alpha}{m}, \frac{\alpha \Psi}{m(\Psi+1)}, (b_q); \end{matrix} x(-\Psi)^m \right] \right\} \quad (65)$$

(vii). Putting  $\beta = \frac{-1}{3}$  and  $\zeta = \Pi$  from equation (31) in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + \frac{2}{3}n)} \binom{\alpha + \frac{2}{3}n}{n}_{p+m} F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) & ; & x \\ \Delta(m; 1 + \alpha - \frac{1}{3}n), (b_q); \end{matrix} \right] t^n = (1 + \Pi)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{3\alpha}{2m}, (a_p); \\ 1 + \frac{3\alpha}{2m}, (b_q); \end{matrix} x(-\Pi)^m \right] \right. \\ \left. + \frac{\mu \Pi}{(1 + \frac{1}{3}\Pi)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{3\alpha}{2m}, 1 + \frac{\alpha \Pi}{m(\Pi+1)}, (a_p); \\ 1 + \frac{3\alpha}{2m}, \frac{\alpha \Pi}{m(\Pi+1)}, (b_q); \end{matrix} x(-\Pi)^m \right] \right\} \quad (66)$$

(viii). Putting  $\beta = \frac{-2}{3}$  and  $\zeta = \Phi$  from equation (32) in equation (33), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + \frac{1}{3}n)} \binom{\alpha + \frac{1}{3}n}{n}_{p+m} F_{q+m} \left[ \begin{matrix} \Delta(m; -n), (a_p) & ; & x \\ \Delta(m; 1 + \alpha - \frac{2}{3}n), (b_q); \end{matrix} \right] t^n = (1 + \Phi)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{3\alpha}{m}, (a_p); \\ 1 + \frac{3\alpha}{m}, (b_q); \end{matrix} x(-\Phi)^m \right] \right. \\ \left. + \frac{\mu \Phi}{(1 + \frac{2}{3}\Phi)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{3\alpha}{m}, 1 + \frac{\alpha \Phi}{m(\Phi+1)}, (a_p); \\ 1 + \frac{3\alpha}{m}, \frac{\alpha \Phi}{m(\Phi+1)}, (b_q); \end{matrix} x(-\Phi)^m \right] \right\} \quad (67)$$

## 6. New Applications of Second Generating Relation (39)

The following Generating relations of this section are believed to be new in author's knowledge and are not available in the literature of Generating relations.

(i). Putting  $\lambda = 1, \mu = \frac{\beta+1}{\alpha}$  in equation (39), using the definition of  $\mathcal{B}_n^{(\alpha, \beta)}(x; m, 1, \frac{\beta+1}{\alpha})$  and after simplifying, we get

$$\sum_{n=0}^{\infty} \binom{\alpha + (\beta + 1)n}{n}_{p+m+1} F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 + \frac{\alpha + (\beta + 1)n}{(\beta + 1)(\beta + 1 - m)}, (a_p); \\ \Delta(m; 1 + \alpha + \beta n), \frac{\alpha + (\beta + 1)n}{(\beta + 1)(\beta + 1 - m)}, (b_q); \end{matrix} x \right] t^n = (1 + \zeta)^\alpha \left\{ \begin{matrix} p+1 F_{q+1} \left[ \begin{matrix} \frac{\alpha}{m(\beta + 1)}, (a_p); \\ 1 + \frac{\alpha}{m(\beta + 1)}, (b_q); \end{matrix} x(-\zeta)^m \right] \\ + \frac{(\beta + 1)\zeta}{(1 - \beta\zeta)} p+2 F_{q+2} \left[ \begin{matrix} \frac{\alpha}{m(\beta + 1)}, 1 + \frac{\alpha\zeta}{(\beta + 1)(1 + m\zeta - \beta\zeta)}, (a_p); \\ 1 + \frac{\alpha}{m(\beta + 1)}, \frac{\alpha\zeta}{(\beta + 1)(1 + m\zeta - \beta\zeta)}, (b_q); \end{matrix} x(-\zeta)^m \right] \end{matrix} \right\} \quad (68)$$

where  $\zeta$  is given by equations (3) and (4).

(ii). Putting  $\lambda = 1, \mu = \frac{\beta+1}{\alpha}, m = 1$  in equation (39) and using the definition of  $\mathcal{B}_n^{(\alpha, \beta)}(x; 1, 1, \frac{\beta+1}{\alpha})$ , we get

$$\sum_{n=0}^{\infty} \binom{\alpha + (\beta + 1)n}{n}_{p+2} F_{q+2} \left[ \begin{matrix} -n, 1 + \frac{\alpha + (\beta + 1)n}{(\beta + 1)\beta}, (a_p); \\ 1 + \alpha + \beta n, \frac{\alpha + (\beta + 1)n}{(\beta + 1)\beta}, (b_q); \end{matrix} x \right] t^n = (1 + \zeta)^\alpha \left\{ \begin{matrix} p+1 F_{q+1} \left[ \begin{matrix} \frac{\alpha}{(\beta + 1)}, (a_p); \\ 1 + \frac{\alpha}{(\beta + 1)}, (b_q); \end{matrix} -x\zeta \right] \\ + \frac{(\beta + 1)\zeta}{(1 - \beta\zeta)} p+2 F_{q+2} \left[ \begin{matrix} \frac{\alpha}{(\beta + 1)}, 1 + \frac{\alpha\zeta}{(\beta + 1)(1 + \zeta - \beta\zeta)}, (a_p); \\ 1 + \frac{\alpha}{(\beta + 1)}, \frac{\alpha\zeta}{(\beta + 1)(1 + \zeta - \beta\zeta)}, (b_q); \end{matrix} -x\zeta \right] \end{matrix} \right\} \quad (69)$$

where  $\zeta$  is given by equations (3) and (4).

(iii). Putting  $\lambda = 1, \mu = \frac{1}{2\alpha}, \beta = \frac{-1}{2}, m = 1$  in equation (39), using the definition of  $\mathcal{B}_n^{(\alpha, -\frac{1}{2})}(x; 1, 1, \frac{1}{2\alpha})$  and replacing  $\zeta$  by  $U$  from equation (29), we get

$$\sum_{n=0}^{\infty} \binom{\alpha + \frac{n}{2}}{n}_{p+2} F_{q+2} \left[ \begin{matrix} -n, 1 - 4\alpha - 2n, (a_p); \\ 1 + \alpha - \frac{n}{2}, -4\alpha - 2n, (b_q); \end{matrix} x \right] t^n = (1 + U)^\alpha \cdot \left\{ \begin{matrix} p+1 F_{q+1} \left[ \begin{matrix} 2\alpha, (a_p); \\ 1 + 2\alpha, (b_q); \end{matrix} -xU \right] \\ + \frac{U}{(2 + U)} p+2 F_{q+2} \left[ \begin{matrix} 2\alpha, 1 + \frac{4\alpha U}{(2 + 3U)}, (a_p); \\ 1 + 2\alpha, \frac{4\alpha U}{(2 + 3U)}, (b_q); \end{matrix} -xU \right] \end{matrix} \right\} \quad (70)$$

(iv). Putting  $\beta = 0$  and  $\zeta = \Theta = \frac{t}{1-t}$  from equation (25) in equation (39), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + n)} \binom{\alpha + n}{n}_{p+m+1} F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 + \frac{\lambda + \mu n}{\mu(1-m)}, (a_p); \\ \Delta(m; 1 + \alpha), \frac{\lambda + \mu n}{\mu(1-m)}, (b_q); \end{matrix} x \right] t^n = (1 - t)^{-\alpha} \left\{ \begin{matrix} \frac{\lambda}{\alpha} p+1 F_{q+1} \left[ \begin{matrix} \frac{\alpha}{m}, (a_p); \\ 1 + \frac{\alpha}{m}, (b_q); \end{matrix} x \left( \frac{-t}{1-t} \right)^m \right] \\ + \frac{\mu t}{(1-t)} p+2 F_{q+2} \left[ \begin{matrix} \frac{\alpha}{m}, 1 + \frac{t\alpha}{\{1+t(m-1)\}}, (a_p); \\ 1 + \frac{\alpha}{m}, \frac{t\alpha}{\{1+t(m-1)\}}, (b_q); \end{matrix} x \left( \frac{-t}{1-t} \right)^m \right] \end{matrix} \right\} \quad (71)$$

(v). Putting  $\beta = 1$  and  $\zeta = \Lambda = \frac{1-2t-\sqrt{(1-4t)}}{2t}$  from equation (26) in equation (39), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + 2n)} \binom{\alpha + 2n}{n}_{p+m+1} F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 + \frac{\lambda + \mu n}{\mu(2-m)}, (a_p); \\ \Delta(m; 1 + \alpha + n), \frac{\lambda + \mu n}{\mu(2-m)}, (b_q); \end{matrix} x \right] t^n = (1 + \Lambda)^\alpha \left\{ \begin{matrix} \frac{\lambda}{\alpha} p+1 F_{q+1} \left[ \begin{matrix} \frac{\alpha}{2m}, (a_p); \\ 1 + \frac{\alpha}{2m}, (b_q); \end{matrix} x(-\Lambda)^m \right] \\ + \frac{\mu\Lambda}{(1-\Lambda)} p+2 F_{q+2} \left[ \begin{matrix} \frac{\alpha}{2m}, 1 + \frac{\Lambda\alpha}{2(1+m\Lambda-\Lambda)}, (a_p); \\ 1 + \frac{\alpha}{2m}, \frac{\Lambda\alpha}{2(1+m\Lambda-\Lambda)}, (b_q); \end{matrix} x(-\Lambda)^m \right] \end{matrix} \right\} \quad (72)$$

(vi). Putting  $\beta = -2$  and  $\zeta = \Xi = \frac{-1+\sqrt{(1+4t)}}{2}$  from equation (27) in equation (39), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha - n)} \binom{\alpha - n}{n}_{p+m+1} F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 - \frac{\lambda + \mu n}{\mu(1+m)}, (a_p); \\ \Delta(m; 1 + \alpha - 2n), \frac{-(\lambda + \mu n)}{\mu(1+m)}, (b_q); \end{matrix} x \right] t^n$$

$$= (1 + \Xi)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} -\frac{\alpha}{m}, (a_p); \\ 1 - \frac{\alpha}{m}, (b_q); \end{matrix} x(-\Xi)^m \right] + \frac{\mu \Xi}{(1 + 2\Xi)^{p+2}} F_{q+2} \left[ \begin{matrix} -\frac{\alpha}{m}, 1 - \frac{\Xi \alpha}{(1+m\Xi+2\Xi)}, (a_p); \\ 1 - \frac{\alpha}{m}, \frac{-\Xi \alpha}{(1+m\Xi+2\Xi)}, (b_q); \end{matrix} x(-\Xi)^m \right] \right\} \quad (73)$$

(vii). Putting  $\beta = -3$  and  $\zeta = \Upsilon$  from equation (28) in equation (39), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha - 2n)} \binom{\alpha - 2n}{n} {}_{p+m+1}F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 - \frac{(\lambda + \mu n)}{\mu(2+m)}, (a_p); \\ \Delta(m; 1 + \alpha - 3n), -\frac{(\lambda + \mu n)}{\mu(2+m)}, (b_q); \end{matrix} x \right] t^n \\ = (1 + \Upsilon)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} -\frac{\alpha}{2m}, (a_p); \\ 1 - \frac{\alpha}{2m}, (b_q); \end{matrix} x(-\Upsilon)^m \right] + \frac{\mu \Upsilon}{(1 + 3\Upsilon)^{p+2}} F_{q+2} \left[ \begin{matrix} -\frac{\alpha}{2m}, 1 - \frac{\alpha \Upsilon}{2(1+m\Upsilon+3\Upsilon)}, (a_p); \\ 1 - \frac{\alpha}{2m}, \frac{-\alpha \Upsilon}{2(1+m\Upsilon+3\Upsilon)}, (b_q); \end{matrix} x(-\Upsilon)^m \right] \right\} \quad (74)$$

(viii). Putting  $\beta = \frac{-1}{2}$  and  $\zeta = U$  from equation (29) in equation (39), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + \frac{1}{2}n)} \binom{\alpha + \frac{1}{2}n}{n} {}_{p+m+1}F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 + \frac{\lambda + \mu n}{\mu(\frac{1}{2}-m)}, (a_p); \\ \Delta(m; 1 + \alpha - \frac{1}{2}n), \frac{\lambda + \mu n}{\mu(\frac{1}{2}-m)}, (b_q); \end{matrix} x \right] t^n \\ = (1 + U)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{2\alpha}{m}, (a_p); \\ 1 + \frac{2\alpha}{m}, (b_q); \end{matrix} x(-U)^m \right] + \frac{\mu U}{(1 + \frac{1}{2}U)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{2\alpha}{m}, 1 + \frac{2U\alpha}{(1+mU+\frac{1}{2}U)}, (a_p); \\ 1 + \frac{2\alpha}{m}, \frac{2U\alpha}{(1+mU+\frac{1}{2}U)}, (b_q); \end{matrix} x(-U)^m \right] \right\} \quad (75)$$

(ix). Putting  $\beta = \frac{-3}{2}$  and  $\zeta = \Psi$  from equation (30) in equation (39), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha - \frac{1}{2}n)} \binom{\alpha - \frac{1}{2}n}{n} {}_{p+m+1}F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 - \frac{(\lambda + \mu n)}{\mu(\frac{1}{2}+m)}, (a_p); \\ \Delta(m; 1 + \alpha - \frac{3}{2}n), -\frac{(\lambda + \mu n)}{\mu(\frac{1}{2}+m)}, (b_q); \end{matrix} x \right] t^n \\ = (1 + \Psi)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} -\frac{2\alpha}{m}, (a_p); \\ 1 - \frac{2\alpha}{m}, (b_q); \end{matrix} x(-\Psi)^m \right] + \frac{\mu \Psi}{(1 + \frac{3}{2}\Psi)^{p+2}} F_{q+2} \left[ \begin{matrix} -\frac{2\alpha}{m}, 1 - \frac{2\Psi\alpha}{(1+m\Psi+\frac{3}{2}\Psi)}, (a_p); \\ 1 - \frac{2\alpha}{m}, \frac{-2\Psi\alpha}{(1+m\Psi+\frac{3}{2}\Psi)}, (b_q); \end{matrix} x(-\Psi)^m \right] \right\} \quad (76)$$

(x). Putting  $\beta = \frac{-1}{3}$  and  $\zeta = \Pi$  from equation (31) in equation (39), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + \frac{2}{3}n)} \binom{\alpha + \frac{2}{3}n}{n} {}_{p+m+1}F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 + \frac{\lambda + \mu n}{\mu(\frac{2}{3}-m)}, (a_p); \\ \Delta(m; 1 + \alpha - \frac{1}{3}n), \frac{\lambda + \mu n}{\mu(\frac{2}{3}-m)}, (b_q); \end{matrix} x \right] t^n \\ = (1 + \Pi)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{3\alpha}{2m}, (a_p); \\ 1 + \frac{3\alpha}{2m}, (b_q); \end{matrix} x(-\Pi)^m \right] + \frac{\mu \Pi}{(1 + \frac{1}{3}\Pi)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{3\alpha}{2m}, 1 + \frac{3\Pi\alpha}{2(1+m\Pi+\frac{1}{3}\Pi)}, (a_p); \\ 1 + \frac{3\alpha}{2m}, \frac{3\Pi\alpha}{2(1+m\Pi+\frac{1}{3}\Pi)}, (b_q); \end{matrix} x(-\Pi)^m \right] \right\} \quad (77)$$

(xi). Putting  $\beta = \frac{-2}{3}$  and  $\zeta = \Phi$  from equation (32) in equation (39), we get

$$\sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{(\alpha + \frac{1}{3}n)} \binom{\alpha + \frac{1}{3}n}{n} {}_{p+m+1}F_{q+m+1} \left[ \begin{matrix} \Delta(m; -n), 1 + \frac{\lambda + \mu n}{\mu(\frac{1}{3}-m)}, (a_p); \\ \Delta(m; 1 + \alpha - \frac{2}{3}n), \frac{\lambda + \mu n}{\mu(\frac{1}{3}-m)}, (b_q); \end{matrix} x \right] t^n \\ = (1 + \Phi)^\alpha \left\{ \frac{\lambda}{\alpha^{p+1}} F_{q+1} \left[ \begin{matrix} \frac{3\alpha}{m}, (a_p); \\ 1 + \frac{3\alpha}{m}, (b_q); \end{matrix} x(-\Phi)^m \right] + \frac{\mu \Phi}{(1 + \frac{2}{3}\Phi)^{p+2}} F_{q+2} \left[ \begin{matrix} \frac{3\alpha}{m}, 1 + \frac{3\Phi\alpha}{(1+m\Phi+\frac{2}{3}\Phi)}, (a_p); \\ 1 + \frac{3\alpha}{m}, \frac{3\Phi\alpha}{(1+m\Phi+\frac{2}{3}\Phi)}, (b_q); \end{matrix} x(-\Phi)^m \right] \right\} \quad (78)$$

Making suitable adjustments of parameters and variables in all generating relations of section 5 and 6, we can also obtain a number of new generating relations involving restricted generalized Laguerre polynomials, restricted Jacobi polynomials, restricted generalized Rice polynomials of Khandekar and other orthogonal polynomials.

## 7. Further Generalizations of Generating relations (33) and (39)

**Generalization of (33):** Let

$$S_n^{(\alpha, \beta)}(x; m) = \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} \binom{\alpha + (\beta + 1)n}{n - mr} \gamma_r x^r \tag{79}$$

where  $\alpha, \beta$  are complex parameters independent of ‘n’;  $m$  is an arbitrary positive integer and  $\{\gamma_r\}$  is a bounded sequence of arbitrary real and complex numbers such that  $\gamma_r \neq 0$ . Then

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{\lambda + \mu n}{\{\alpha + (\beta + 1)n\}} S_n^{(\alpha, \beta)}(x; m) t^n &= (1 + \zeta)^\alpha \left\{ \lambda \sum_{n=0}^{\infty} \frac{1}{\{\alpha + (\beta + 1)m n\}} \gamma_n x^n \zeta^{m n} \right. \\ &\quad \left. + \frac{\mu}{(1 - \beta \zeta)} \sum_{n=0}^{\infty} \frac{\{\zeta \alpha + m(1 + \zeta)n\}}{\{\alpha + (\beta + 1)m n\}} \gamma_n x^n \zeta^{m n} \right\} \end{aligned} \tag{80}$$

where  $\zeta$  is given by

$$\zeta = t(1 + \zeta)^{(\beta + 1)}; \zeta(0) = 0 \tag{81}$$

provided that each of the series involved is absolutely convergent.

**Independent Demonstration:** Using the definition (79) of  $S_n^{(\alpha, \beta)}(x; m)$  in left hand side of equation (80), we get

$$\begin{aligned} \Omega^{**} &= \sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{\{\alpha + (\beta + 1)n\}} S_n^{(\alpha, \beta)}(x; m) t^n \\ &= \sum_{n=0}^{\infty} \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} \frac{(\lambda + \mu n) \Gamma\{\alpha + (\beta + 1)n + 1\}}{\{\alpha + (\beta + 1)n\} \Gamma(n - mr + 1) \Gamma\{\alpha + \beta n + mr + 1\}} \gamma_r x^r t^n \end{aligned} \tag{82}$$

Applying summation identity (10) and then simplifying further, we get

$$\Omega^{**} = \sum_{r=0}^{\infty} \frac{\gamma_r x^r (t)^{mr}}{\{\alpha + m(\beta + 1)r\}} \left\{ \sum_{n=0}^{\infty} \frac{\{\alpha + (\beta + 1)mr\}(\lambda + \mu n + \mu mr)}{\{\alpha + (\beta + 1)mr + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)mr + (\beta + 1)n}{n} t^n \right\} \tag{83}$$

Now using first modified Gould’s identity (7) with condition (81), we get

$$\Omega^{**} = (1 + \zeta)^\alpha \sum_{r=0}^{\infty} \left[ \lambda + \left\{ \mu mr + \frac{\mu \zeta \{\alpha + (\beta + 1)mr\}}{(1 - \beta \zeta)} \right\} \right] \frac{\gamma_r x^r (\zeta)^{mr}}{\{\alpha + m(\beta + 1)r\}} \tag{84}$$

Changing the summation index from  $r$  to  $n$  and after solving it further, we get the general result (80) corresponding to our first generating relation (33) subject to the conditions (81).

**Generalization of (39):** Let

$$T_n^{(\alpha, \beta)}(x; m, \lambda, \mu) = \frac{1}{(\lambda + \mu n)} \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} \left\{ \binom{\alpha + (\beta + 1)n}{n - mr} [\lambda + \mu n + \mu(\beta + 1 - m)r] \right\} \gamma_r x^r \tag{85}$$

where  $\alpha, \beta, \lambda, \mu$  are complex parameters independent of ‘n’;  $m$  is an arbitrary positive integer and  $\{\gamma_r\}$  is a bounded sequence of arbitrary real and complex numbers such that  $\gamma_r \neq 0$ . Then

$$\begin{aligned} \sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{\{\alpha + (\beta + 1)n\}} T_n^{(\alpha, \beta)}(x; m, \lambda, \mu) t^n &= (1 + \zeta)^\alpha \left\{ \sum_{n=0}^{\infty} \frac{\lambda}{\{\alpha + (\beta + 1)m n\}} \gamma_n x^n \zeta^{m n} \right. \\ &\quad \left. + \frac{\mu}{(1 - \beta \zeta)} \sum_{n=0}^{\infty} \frac{1}{\{\alpha + (\beta + 1)m n\}} \gamma_n x^n \zeta^{m n} \right\} \end{aligned} \tag{86}$$

where  $\zeta$  is given by  $\zeta = t(1 + \zeta)^{(\beta+1)}$ ;  $\zeta(0) = 0$  provided that each of the series involved is absolutely convergent.

**Independent Demonstration:** Using the definition (85) of  $T_n^{(\alpha,\beta)}(x; m, \lambda, \mu)$  in left hand side of equation (86), we get

$$\begin{aligned}\Omega^{***} &= \sum_{n=0}^{\infty} \frac{(\lambda + \mu n)}{\{\alpha + (\beta + 1)n\}} T_n^{(\alpha,\beta)}(x; m, \lambda, \mu) t^n \\ &= \sum_{n=0}^{\infty} \sum_{r=0}^{\lfloor \frac{n}{m} \rfloor} \left\{ \binom{\alpha + (\beta + 1)n}{n - mr} \frac{\{\lambda + \mu n + \mu(\beta + 1)r - \mu mr\}}{\{\alpha + (\beta + 1)n\}} \right\} \gamma_r x^r t^n\end{aligned}\quad (87)$$

Applying summation identity (10) and then simplifying further, we get

$$\Omega^{***} = \sum_{r=0}^{\infty} \frac{\gamma_r x^r (t)^{mr}}{\{\alpha + m(\beta + 1)r\}} \left\{ \sum_{n=0}^{\infty} \frac{\{\alpha + (\beta + 1)mr\} \{\lambda + \mu n + \mu(\beta + 1)r\}}{\{\alpha + (\beta + 1)mr + (\beta + 1)n\}} \binom{\alpha + (\beta + 1)mr + (\beta + 1)n}{n} t^n \right\} \quad (88)$$

Now using second modified Gould's identity (8) with conditions (81), we get

$$\Omega^{***} = (1 + \zeta)^\alpha \sum_{r=0}^{\infty} \left[ \lambda + \left\{ \mu(\beta + 1)r + \frac{\mu\zeta \{\alpha + (\beta + 1)mr\}}{(1 - \beta\zeta)} \right\} \right] \cdot \frac{\gamma_r x^r (\zeta)^{mr}}{\{\alpha + m(\beta + 1)r\}} \quad (89)$$

Changing the summation index from  $r$  to  $n$  and after solving it further, we get the general result (86) corresponding to our second generating relation (39) subject to the conditions (81). In the definitions of generalized polynomials given by  $S_n^{(\alpha,\beta)}(x; m)$  and  $T_n^{(\alpha,\beta)}(x; m, \lambda, \mu)$ , putting

$$\gamma_r = \frac{(-1)^{mr} (a_1)_r \dots (a_p)_r}{(b_1)_r \dots (b_q)_r r!},$$

we obtain Srivastava's generalized hypergeometric polynomials of one variable  $H_n^{(\alpha,\beta)}(x; m)$  and Pathan's generalized hypergeometric polynomials of one variable  $\mathcal{B}_n^{(\alpha,\beta)}(x; m, \lambda, \mu)$  respectively.

## References

- [1] J.W.Brown, *On Zero type sets of Laguerre polynomials*, Duke Math. J., 35(1968), 821-823.
- [2] J.W.Brown, *New generating functions for classical polynomials*, Proc. Amer. Math. Soc., 21(1969), 263-268.
- [3] J.W.Brown, *New Generating Functions for Classical Polynomials(Abstract)*, Notices Amer. Math. Soc., 16(1969), 412.
- [4] J.W.Brown, *On a Generating Function for Laguerre Polynomials*, Ricerca(Napoli), (2)(1970), 26-28.
- [5] L.Carlitz, *Some Generating Functions for Laguerre Polynomials*, Duke Math. J., 35(1968), 825-827.
- [6] H.W.Gould, *Some generalizations of Vandermonde's Convolution*, Amer. Math. Monthly, 63(1956), 84-91.
- [7] H.W.Gould, *A Series Transformation for Finding Convolution Identities*, Duke. Math. J., 28(1961), 193-202.
- [8] C.M.Joshi and M.L.Prajapat, *A Triple integral transformation and its applications to generating functions*, Boll. Un. Mat. Ital., (5)14A(1977), 264-274.
- [9] P.R.Khandekar, *On a generalization of Rice's polynomials*, I. Proc. Nat. Acad. Sci. India Sect. A34(1964), 157-162.
- [10] G.Pólya and G.Szegő, *Problems and Theorems in Analysis*, Vol.I, (Translated from the German by D.Aeppli), Springer-Verlag, New York, Heidelberg and Berlin, (1972).
- [11] E.D.Rainville, *Special Functions*, The Macmillan, New York, (1960); Reprinted by Chelsea Publishing Company, Bronx, New York, (1971).
- [12] J.Riordan, *Combinatorial Identities*, John Wiley & Sons, New York, London and Sydney, (1968).
- [13] B.L.Sharma and R.F.A.Abiodun, *A new Generating Function for a generalized function of two variables*, Proc. Amer. Math. Soc., 53(1975), 379-384.

- [14] R.M.Shreshtha, *Generating Functions for generalized Hypergeometric series*, C.R. Acad. Bulgare Sci., 31(1978), 389-391.
- [15] H.M.Srivastava, *Generating Functions for Jacobi and Laguerre Polynomials*, Proc. Amer. Math. Soc., 23(1969), 590-595.
- [16] H.M.Srivastava, *A class of Generating Functions for generalized Hypergeometric Polynomials(Abstract)*, Notices Amer. Math. Soc., 16(1969), 975.
- [17] H.M.Srivastava, *A class of Generating Functions for generalized Hypergeometric Polynomials*, J. Math. Anal. Appl., 35(1971), 230-235.
- [18] H.M.Srivastava, *A Note on Certain Generating functions for the Classical Polynomials*, Atti Accad. Naz. Lincei Rend. Cl. Sci.Fis. Mat. Natur., 63(8)(1977), 328-333.
- [19] H.M.Srivastava, *Some Generating functions for Laguerre and Bessel Polynomials*, Bull. Inst. Math. Acad. Sinica, 8(1980), 571-579.
- [20] H.M.Srivastava and H.L.Manocha, *A Treatise on Generating functions*, Halsted Press (Ellis Horwood Ltd., Chichester, U.K.), John Wiley and Sons, New York, Chichester, Brisbane and Toronto, (1984).
- [21] E.T.Whittaker and G.N.Watson, *A Course of Modern Analysis*, Fourth ed., Cambridge Univ.Press, Cambridge, London and New York, (1927).
- [22] D.Zeitlin, *A new class of Generating Functions for Hypergeometric Polynomials*, Proc. Amer. Math. Soc., 25(1970), 405-412.